

# **WHITE MATTER CORRELATES OF NEUROPSYCHOLOGICAL FUNCTION IN YOUNG ADULT METHAMPHETAMINE USERS**

**Dr. Carla Patricia Freeman**

**MBChB (UCT) FC Psych (SA) MMed Psych (UCT)**

**Student number: FRMCAR002**

Thesis presented for the partial fulfillment of the degree of Master of  
Philosophy in Neuropsychiatry in the Division of Neuropsychiatry, Department  
of Psychiatry and Mental Health, University of Cape Town

JUNE 2016

Supervisor: Associate Professor John A Joska<sup>1</sup>

Co-supervisor: Professor Dan J Stein<sup>2</sup>

1. Head of Division of Neuropsychiatry, Department of Psychiatry and Mental Health, University of Cape Town

2. Head of Department of Psychiatry and Mental Health, University of Cape Town

The copyright of this thesis vests in the author. No quotation from it or information derived from it is to be published without full acknowledgement of the source. The thesis is to be used for private study or non-commercial research purposes only.

Published by the University of Cape Town (UCT) in terms of the non-exclusive license granted to UCT by the author.

## DECLARATION

I, Carla Patricia Freeman, hereby declare that the work on which this dissertation is based is my original work (except where acknowledgements indicate otherwise) and that neither the whole or any part has been, is being, or is to be submitted for any other degree at this or any other university.

I empower the University of Cape Town to reproduce for the purpose of research either the whole or any portion of the contents in any manner whatsoever.

Signature:

Signed by candidate

Signature removed

Signed at: Cape Town on the 9<sup>th</sup> day of June 2016.

## PLAGIARISM DECLARATION FOR WRITTEN WORK

I, **Dr Carla Patricia Freeman**, as a candidate registered for an MPhil Neuropsychiatry degree at the University of Cape Town declare the following:

1. I am aware that plagiarism is the use of someone else's work without their permission and/or without acknowledging the original source.
2. I am aware that plagiarism is wrong.
3. I confirm that the written work is my own unless stated otherwise.
4. I have followed the required conventions in referencing the thoughts and ideas of others.
5. I understand that the University of Cape Town may take disciplinary action against me if there is a belief that this is not my own original work or if I have failed to acknowledge the ideas or writing of others.

Signature:

Signed by candidate

signature removed

Signed at: Cape Town on the 9<sup>th</sup> day of June 2016.

## **ACKNOWLEDGMENTS**

A number of individuals have contributed to the success of this research study. Firstly, I wish to thank both of my supervisors, Professors Joska and Stein, for their contribution to this manuscript as well as their mentorship throughout my training. Secondly, to the participants willing to dedicate both their time and experience to this study. Thirdly to my colleagues: Jean Paul Fouche; Jonathan Ipser and Hetta Gouse, all of whom have whole-heartedly contributed to this research study and, finally, to the Discovery Foundation for allowing me the opportunity to undertake this research.

This manuscript is dedicated to my brother, Robin, who taught me much as he succumbed to his battle with addiction.

## **ABSTRACT**

### *Background*

Methamphetamine (MA) abuse is a global health concern due to widespread use and harmful effects, which includes neurotoxicity. This study aimed to describe neurocognitive deficits associated with MA dependence in young adults and to explore whether these deficits correlate with white matter (WM) microstructural abnormalities using diffusion tensor imaging (DTI).

### *Methods*

Twenty-one MA dependent individuals recently enrolled in an outpatient rehabilitation program and nineteen healthy controls participated in the study. Each participant completed a neuropsychological evaluation and underwent diffusion tensor imaging within one week of testing. Average whole-brain fractional anisotropy (FA) and mean diffusion (MD) measures derived from DTI data were compared between groups. Group differences in performance within specific neurocognitive domains and in a composite global neurocognitive score (GNS) were tested using non-parametric univariate statistics and within a linear regression framework, adjusting for age and gender. Correlation analyses were conducted to test associations between the neuropsychological data and selected frontal white matter (WM) tracts, including the genu and body of the corpus callosum (CC); right and left cingulum bundle (CB); right and left uncinate fasciculus (UF); right and left anterior corona radiata (CR) and the right and left superior longitudinal fasciculus (SLF).

### *Results*

No significant between-group differences were detected for performance in any of the neuropsychological domains assessed. No relationship between FA or MD and the GNS was demonstrated in the tracts of interest. After adjusting for age and gender, significant group differences in FA and MD were detected across several regions of interest (ROI), however, these did not survive corrections for multiple comparisons.

### *Conclusion*

Cognitive performance and white matter integrity did not differ between young MA dependent subjects and healthy controls. Whatever differences that were found in white matter did not survive correction for multiple comparisons. These findings may reflect one or more of several possibilities: that brain function and structure is relatively preserved in younger individuals; or that differences were too small to be detected in this sample. Further studies should explore the effects of aging, poly-substance abuse and HIV co-infection on neurocognitive functioning and structural brain integrity in methamphetamine users.

## TABLE OF CONTENTS

DECLARATION.....	2
PLAGIARISM DECLARATION FOR WRITTEN WORK.....	3
ACKNOWLEDGEMENTS.....	4
ABSTRACT.....	5
TABLE OF CONTENTS.....	7
LIST OF ABBREVIATIONS.....	9
CHAPTER 1: INTRODUCTION.....	10
1.1    Context.....	10
1.1.1  Methamphetamine use in South Africa.....	10
1.1.2  Methamphetamine and neurotoxicity.....	10
1.1.3  Methamphetamine and cognition.....	11
1.1.4  Methamphetamine and diffusion tensor imaging.....	12
1.1.5  Literature review of DTI studies in methamphetamine dependence.....	13
1.2    Ethical considerations.....	15
1.3    Author Guidelines for PSYCHIATRY RESEARCH: NEUROIMAGING.....	17
1.4    References.....	18
CHAPTER 2: PUBLICATION READY FORMAT.....	21
2.1    White matter correlates of neuropsychological function in abstinent, young methamphetamine users – a diffusion tensor imaging study.....	
2.2    References.....	34
2.3    List of tables:	
2.3.1  Table 1: Participant demographic data and clinical characteristics.....	37
2.3.2  Table 2: Neuropsychological performance of MA users and healthy comparison subjects.....	38
2.3.3  Table 3: Differences between MA+ and MA- groups on DTI outcomes for tracts of interest.....	39
2.3.4  Table 4: DTI correlates of Global Neurocognitive Score .....	40



## SUPPLEMENTARY MATERIAL

- Table S1: Mean DTI values in MA dependent individuals and healthy control subjects .....41
- Table S2: Adjusted group differences in DTI metrics for fractional anisotropy (FA) .....42
- Table S3: Adjusted group differences in DTI metrics for mean diffusion (MD) .....44

## APPENDIX

APPENDIX A: HREC approval letter 263/2010 .....	47
APPENDIX B: HREC approval letter 635/2014.....	48
APPENDIX C: MATRIX information session (slides): 28 June 2011 .....	49
APPENDIX D: Candidate approval for MPhil (Neuropsychiatry) UCT .....	52
APPENDIX E: City of Cape Town permission to recruit from Matrix Clinic ...	53
APPENDIX F: Inclusion/exclusion criteria flyer for Matrix and Mitchells Plain Community Health Centre .....	54
APPENDIX G: Participant informed consent document .....	55
APPENDIX H: Author guidelines Psychiatry Research: Neuroimaging .....	57

## ABBREVIATIONS

(CUBIC)      Cape Universities Brain Imaging Centre

(CB)	Cingulum Bundle
(CC)	Corpus Callosum
(CR)	Corona Radiata
(DAT)	Dopamine transporter
(DTI)	Diffusion tensor imaging
(FA)	Fractional anisotropy
(FSL)	FMRIB Software Library
(GNS)	Global Neurocognitive Score
(HREC)	Human Research Ethics Committee
(MA)	Methamphetamine
(MA +)	Methamphetamine user
(MA -)	Healthy control subject
(MD)	Mean diffusion
(MRI)	Magnetic resonance imaging
(PET)	Positron emission tomography
(ROI)	Region of interest
(SA)	South Africa
(SACENDU)	South African Community Epidemiology Network on Drug Use
(SLF)	Superior Longitudinal Fasciculus
(5-HT)	Serotonin
(TBSS)	Tract-based spatial statistics
(UCT)	University of Cape Town
(UDS)	Urine Drug Screen
(UF)	Uncinate Fasciculus
(WC)	Western Cape
(WM)	White matter

## **CHAPTER 1: INTRODUCTION**

### **1.1 Context**

#### **1.1.1 Methamphetamine use in South Africa**

Methamphetamine (MA), like cocaine, is a powerful stimulant drug of abuse with widespread illegal use. It's popularity peaked in the United States (US) in the 1960s and later in the 1990s, with production largely driven by small-scale informal laboratories making use of readily available ingredients, e.g. pseudoephedrine in cold and flu medication (Ciccarone 2011). MA abuse is traditionally associated with certain subgroups (e.g. gang members; sex workers; truck drivers; gay men), although growing concern has been expressed over its escalating use in South Africa (SA), particularly the Western Cape population (Brecht et al. 2004; Pluddemann et al. 2010). Despite numerous negative consequences affecting social, physical, psychological and occupational well-being, South Africans continue to abuse MA. Furthermore, in the context of South Africa's high HIV prevalence, the associated risk of HIV exposure, increased transmission rates, treatment non-adherence and disease progression in individuals who abuse MA is of grave concern (Kapp 2009; Colfax et al. 2010; Carrico et al. 2011). This chapter will outline some of the known deleterious effects of MA on human brain structure and function and will further describe the ethical considerations necessary for research in this study population.

#### **1.1.2 Methamphetamine and neurotoxicity**

MA is a highly lipophilic, neurotoxic molecule capable of producing significant disruption in brain structure and function. Animal and human studies support a dose-related association between acute and chronic MA use and neurotoxicity. Rats exposed to MA exhibit decreased levels of striatal dopamine as well as destruction of dopaminergic nerve terminals (Ricaurte et al. 1982). Serotonergic function is also compromised following MA exposure as evidenced by: loss of the 5-HT transporter; 5-HT depletion; depletion of 5-

HT metabolites and reductions in tryptophan hydroxylase (as reviewed in Meredith et al 2005) (Meredith et al. 2005). Post mortem findings further support alterations in dopaminergic functioning seen in neuroimaging studies as evidenced by deficits in available dopamine, the dopamine transporter and tyrosine hydroxylase, a dopamine precursor (Wilson et al. 1996).

Proposed mechanisms of neurotoxicity include accumulation of reactive oxygen species and severe oxidative stress, glutamate mediated excitotoxicity and mitochondrial dysfunction, preferentially affecting striatal dopaminergic nerve terminals (Quinton & Yamamoto 2006). Davidson et al. further postulate that MA induced neurotoxicity occurs via two pathways i.e. loss of dopaminergic function in acute overdose (4 x 5mg/kg) and MA induced apoptosis seen in chronic use (15mg/kg/day x 14 days) (Davidson et al. 2001).

### **1.1.3 Methamphetamine and cognition**

Long-term/chronic MA use has been shown to result in significant cognitive impairment affecting multiple domains. Up to 40% of long-term MA dependent individuals exhibit global neuropsychological impairment (Rippeth et al. 2004). Cognitive processes dependent on frontal-striatal and limbic circuits are predominantly affected in MA abuse, irrespective of age and level of education (Scott et al. 2007). The following domains have been shown to be impaired: episodic memory; prospective memory; executive function, complex information processing speed and psychomotor function (Scott et al. 2007; Rendell et al. 2008). Smaller effect sizes were noted in attention/working memory, language and visual construction. Successful rehabilitation, including therapeutic strategies, abstinence and impulse control, rely heavily on the aforementioned neuropsychological domains being intact. Impairment in areas such as learning, memory and impulse control may help to explain the difficulty experienced by MA users, engaged in a rehabilitation program, to refrain from using the drug (Gowin et al. 2013).

Despite increasing numbers of individuals using MA, few studies of this nature exist in South African populations. This research may further enhance our understanding of the complex maladaptive behaviours seen in substance-dependent individuals.

#### **1.1.4 Methamphetamine and diffusion tensor imaging**

Magnetic resonance imaging (MRI) plays an increasingly important role in understanding brain structure and function in both normal and diseased states. Diffusion tensor imaging (DTI) was introduced in 1994 and is one of several MRI modalities that allow for the non-invasive study of white matter integrity based on the free diffusion of water molecules within white matter tracts (Assaf & Pasternak 2007; Le Bihan et al. 2001). As reviewed in Salo and Fassbender, a rich body of literature of animal studies exists supporting the deleterious effects of MA on brain white matter, however, few human studies have been published (Tabled below) (Salo & Fassbender 2011). Furthermore, the combination of DTI parameters correlated with cognitive measures could provide a useful marker of human behaviour and response, for example, predicting treatment outcomes in a rehabilitation setting.

DTI employs the principle of Brownian motion i.e. the random movement of water molecules within a space. This motion is deemed isotropic if movement occurs equally in all directions. If water molecules are restricted in a plane e.g. along the axis of a neuron, movement will occur at different speeds with those along the axon moving the fastest compared to molecules moving perpendicular to the axis. This property of molecules in motion is called anisotropy and, with the aid of sophisticated MRI techniques, can be used to demonstrate the direction of water molecules along white matter tracts. These in turn allow clinicians to visualize anisotropic diffusion of water molecules within several cubic mm of white matter, known as a voxel, which may be representative of several disease states (Le Bihan et al. 2001). The resultant anisotropy allows for information to be gained on both tissue microstructure and architecture within the voxel. DTI has been successfully employed to

demonstrate disruption of brain tissue in a number of pathological states. Examples include: demyelination in multiple sclerosis and oedema in response to inflammation seen in acute cerebrovascular accidents (CVA) (Filippi et al. 2001; Sotak 2002).

Fractional anisotropy (FA) is the most frequently used parameter of DTI. FA provides information on the orientation and integrity of white matter fibers within a voxel. Each voxel consists of thousands of white matter fibers and a supportive matrix of glial cells, including oligodendrocytes responsible for the myelin sheath around the axon. FA comprises of an intensity scale that ranges between zero and one. FA is high (close to one) if diffusivity is rapid along fibres and approaches zero if diffusion occurs in all directions (Assaf & Pasternak 2007). Mean diffusivity (MD) provides an overall measure of diffusivity within a voxel or region. MD is understood to increase in pathological disease states e.g. inflammation and oedema (Le Bihan et al. 2001). FA is thought to decrease in the presences of tissue damage, due to either increased radial (RD) or decreased axial diffusivity (AD). More recently, imaging specialists have started to examine AD and RD as separate entities, in combination with FA, in order to glean more information on tissue structure (Alexander et al. 2007). Several limitations should be taken into consideration when interpreting data obtained using DTI. These include: artifact; thermal and physiological image noise; dilution of volume when averaged in larger voxels and fibre crossing (Alexander et al. 2007).

#### **1.1.5 Literature review of DTI studies in methamphetamine dependence**

A literature review conducted, using Pubmed Central, of DTI studies undertaken in MA dependence yielded the results summarized in Table X below. Keywords used in the search include: 'DTI', 'diffusion tensor imaging' 'methamphetamine', 'MA', 'amphetamine' and 'structural imaging' (Alicata et al. 2009; Chung et al. 2007; Ersche et al. 2012; Kim et al. 2009; Salo et al. 2009; Tobias et al. 2010; Lederer et al. 2015).

**Table I: Literature review of DTI studies in MA dependence**

Primary author	Year	Sample	Whole brain voxel analysis or ROI	Measure	Significant findings
Alicata	2009	30 MA +, 30 MA -; age 33.3 (8.6)	Frontal WM; parietal WM; genu CC; caudate; putamen; thalamus; cerebellar vermis	FA, ADC	MA +: decreased FA in right frontal WM; increased ADC in left caudate and bilateral putamen. Higher ADC in left putamen and higher right putamen was associated with higher MA doses.
Chung	2006	32 MA +, 30 healthy controls; men (n=23), women (n-9); age men 36.0 (6.7), women 29.0 (7.2)	Bilateral frontal WM; bilateral parietal WM; bilateral occipital WM. (frontal matter measured relative to the AC-PC plane	FA	MA +: Decreased FA in bilateral frontal WM at the AC-PC plane and right frontal WM above this plane. Less prominent in female participants.
Ersche	2012	50 sibling pairs with at least 1 sibling MA+, 50 unmatched controls	Tract-based skeleton	FA	Decreased mean FA in MA + and their siblings compared to healthy controls. Therefore observed WM abnormalities were shared among family members and may have pre-disposed them to drug-taking.
Kim	2008	11 MA + abstinent, 13 healthy controls (Not tested for HIV); age 34.4 (2.9)	Genu and splenium of the CC	FA	Decreased FA in genu; trend to decreased FA in splenium although not significant.
Lederer	2016	40 MA +, 40 matched controls	Genu CC, Cingulum, Uncinate fasciculus	FA, MD, parallel diffusivity, perpendicular diffusivity	Increased MD and increased perpendicular diffusivity in the genu of the corpus callosum. Increased MD in the right cingulum. None of the WM changes were significantly associated with aggression.

Salo	2009	37 MA +, 17 healthy controls	Genu and splenium of the CC	FA, ADC	No significant difference in FA and ADC across groups. Trend towards decreased FA in genu but not significant (p=0.9)
Tobias	2010	23 MA +, 18 healthy controls	1) Prefrontal WM: four levels inferior-superior to the AC-PC plane (as per Chung, 2007) 2) Midline genu; midline splenium; bilateral anterior, midcaudal and caudal superior CR; anterior and posterior limbs and genu of the internal capsule; and the perforant path of the hippocampal formation.	FA	Lower FA: Right prefrontal WM above the AC-PC; midline genu CC; left and right midcaudal superior corona radiate and right perforant fibres.

WM = White matter; AC-PC = Anterior commissure-posterior commissure

CR = Corona radiata

Few studies appear to have been conducted in this area despite many authors emphasizing the value of DTI in understanding microstructural abnormalities associated with MA addiction. Replication of these studies in a South African, MA dependent population, correlating neuropsychological deficits with white matter deficits seen on DTI, will be of great value and may provide further insight into understanding maladaptive behavioural patterns associated with MA abuse.

## 1.2 Ethical considerations



Good clinical practice principles ensure that research conducted in clinical populations adheres to sound ethical and scientific standards, at all times, in order to protect the rights, safety and well being of study participants. Several special considerations need to occur when research is conducted with human subjects who abuse substances. Firstly, ethical approval of the study protocol, in accordance with the principles laid down by the Declaration of Helsinki, must occur prior to the commencement of research. This research study was approved by the Human Research Ethics Committee (HREC) of the University of Cape Town on the 6<sup>th</sup> of July 2010 (HREC 263/2010), initially as part of a larger parent study, i.e. Methamphetamine induced deficits in neuro-and-social cognition in the context of HIV infection (Appendix A), and later as a separate protocol for the achievement of a Masters of Philosophy (MPhil) in Neuropsychiatry on the 29<sup>th</sup> of August 2014 [HREC 635/2014] (Appendix B). For the purpose of fulfillment of the MPhil requirements, this thesis focused exclusively on HIV negative, methamphetamine dependent individuals and HIV negative, healthy controls. Permission was obtained from both the City of Cape Town and the Provincial Government of the Western Cape to conduct research in the relevant treatment facilities (Appendix E).

According to the Belmont Report, the following ethical principles should guide all clinical research: respect for persons; beneficence and justice. I will address each principle individually with regards to this research study.

(i) Respect for persons: every participant was comprehensively informed of the study prior to enrollment. I presented an information session to both staff and incumbents at Matrix clinic on the known effects of MA on the brain to ensure adequate knowledge on the purpose of the research (Appendix C). Participation was voluntary and all subjects were informed that they were able to withdraw from the study at any point without negative consequences. All participants were actively enrolled in their rehabilitation programme and had full capacity, thus ensuring competence to sign informed consent. Acutely intoxicated participants were not considered for the study. The investigators covered all transport costs and participants were compensated with food vouchers for the two days spent engaged in study procedure. Participants were not compensated with cash at any point due to the risk of this being

used to purchase illicit drugs. Participants were free of undue influence and not coerced to participate in the study. (ii) Beneficence: the investigators maintained ongoing awareness of minimizing risk for participants while maximizing potential benefit without permanent injury at all times. Due to the cross-sectional nature of the study, potential risks mostly pertained to procedure on the study day. Study participants were informed of all study procedures i.e. neuromedical exam; neuropsychological testing and MRI, with their comfort and safety prioritized at all times. (iii) Justice: selection of study participants ensured fair distribution of the risks and benefits of the research. Members of the healthy control group comprised of family and friends of the MA dependent group and were invested in advancing scientific knowledge of MA abuse despite not being addicts themselves.

In addition to the above considerations, the counselors at Matrix were informed of those participating in this research study. With the participants consent, essential clinical information was shared with the counselor to ensure adequate support for the participant throughout the study process.

### **1.3 Author Guidelines for PSYCHIATRY RESEARCH: NEUROIMAGING**

The journal Psychiatry Research is the official publication of The International Society for Neuroimaging in Psychiatry. The Neuroimaging section caters for a variety of publications related to brain imaging of psychiatric disorders including: structural imaging; effects of behavioural tasks and impact of medication within this realm. The journal carries an impact factor of 2.831 (© Thomas Reuters Journal Citation Reports 2014).

The selection of this journal is an appropriate platform for my study as it encompasses brain imaging of a known noxious agent to the brain (methamphetamine) and correlates these findings with cognitive deficits present in this study population. These cognitive deficits in turn are associated with certain deleterious behaviours e.g. executive dysfunction resulting in poor

planning and execution of rehabilitation endeavours. The findings of such a study may help to link structural imaging with associated human behaviours, furthering our scientific understanding of addiction and subsequent treatment planning. This journal is listed on the Thomas Reuters/Institute for Scientific Information (ISI) Web of Science list January 2014: ISSN 0925-4927; E-ISSN 1872-7506; Netherlands.

#### 1.4 REFERENCES: CHAPTER 1

- Alexander, A. et al., 2007. Diffusion tensor imaging of the brain. *Neurotherapeutics*, 4(3), pp.316–329.
- Alicata, D. et al., 2009. Higher diffusion in striatum and lower fractional anisotropy in white matter of methamphetamine users. *Psychiatry Research: Neuroimaging*, 174, pp.1–8.
- Assaf, Y. & Pasternak, O., 2007. Diffusion Tensor Imaging (DTI)-based White Matter Mapping in Brain Research: A Review. *J Mol Neurosci*, 34, pp.51–61.
- Brecht, M.-L. et al., 2004. Methamphetamine use behaviors and gender differences. *Addictive Behaviors*, 29(1), pp.89–106.
- Carrico, A.W. et al., 2011. Psychiatric Risk Factors for HIV Disease Progression: The Role of Inconsistent Patterns of Antiretroviral Therapy Utilization. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 56(2), pp.146–150.
- Chung, A. et al., 2007. Decreased frontal white-matter integrity in abstinent methamphetamine abusers. *International Journal of Neuropsychopharmacology*, 10, pp.765–775.
- Ciccarone, D., 2011. Stimulant Abuse: Pharmacology, Cocaine, Methamphetamine, Treatment, Attempts at Pharmacotherapy. *Primary Care: Clinics in Office Practice*, 38(1), pp.41–58. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3056348/pdf/nihms264606.pdf>.
- Colfax, G. et al., 2010. HIV in people who use drugs 5Amphetamine-group substances and HIV. *The Lancet*, 376(9739), pp.458–474.
- Davidson, C. et al., 2001. Methamphetamine neurotoxicity: necrotic and apoptotic mechanisms and relevance to human abuse and treatment. *Brain Research Reviews*, 36, pp.1–22.

- Ersche, K.D. et al., 2012. Abnormal Brain Structure Implicated in Stimulant Drug Addiction. *Science*, 335, pp.601–604.
- Filippi, M. et al., 2001. Diffusion tensor magnetic resonance imaging in multiple sclerosis. *Neurology*, 56, pp.304–311.
- Gowin, J.L. et al., 2013. Altered cingulate and insular cortex activation during risk-taking in methamphetamine dependence: losses lose impact. *Addiction*, 109(2), pp.237–247.
- Kapp, C., 2009. Crystal meth boom adds to South Africa's health challenges. *Lancet*, 371, pp.193–194.
- Kim, I.-S. et al., 2009. Reduced corpus callosum white matter microstructural integrity revealed by diffusion tensor eigenvalues in abstinent methamphetamine addicts. *Neurotoxicology*, 30, pp.209–213.
- Le Bihan, D. et al., 2001. Diffusion Tensor Imaging: Concepts and Applications. *JOURNAL OF MAGNETIC RESONANCE IMAGING*, 13, pp.534–546.
- Lederer, K. et al., 2015. Frontal white matter changes and aggression in methamphetamine dependence. *Metabolic Brain Disease*, 31(1), pp.53–62.
- Meredith, C.W. et al., 2005. Implications of Chronic Methamphetamine Use: A Literature Review. *Harv Rev Psychiatry*, pp.1–14.
- Pluddemann, A. et al., 2010. Monitoring Alcohol & Drug Abuse Trends in South Africa. *SACENDU Research Brief*, 13(2), pp.1–16.
- Quinton, M.S. & Yamamoto, B.K., 2006. Causes and Consequences of Methamphetamine and MDMA Toxicity. *AAPS Journal*, 8(2), pp.337–347.
- Rendell, P.G., Mazur, M. & Henry, J.D., 2008. Prospective memory impairment in former users of methamphetamine. *Psychopharmacology*, 203(3), pp.609–616.
- Ricaurte, G.A. et al., 1982. Dopamine nerve terminal degeneration produced by high doses of methylamphetamine in the rat brain. *Brain Res.*, 235, pp.93–103.
- Rippeth, J.D. et al., 2004. Methamphetamine dependence increases risk of neuropsychological impairment in HIV infected persons. *Journal of International Neuropsychological Society*, 10, pp.1–14.
- Salo, R. & Fassbender, C., 2011. Structural, Functional and Spectroscopic MRI Studies of Methamphetamine Addiction. In *Current Topics in Behavioral Neurosciences*. Current Topics in Behavioral Neurosciences. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 321–364. Available at: <http://www.springerlink.com/content/q61067r133576278/fulltext.pdf>.

- Salo, R. et al., 2009. Cognitive Control and White Matter Callosal Microstructure in Methamphetamine Dependent Subjects: A DTI Study. *Biological Psychiatry*, 65(2), pp.122–128.
- Scott, J.C. et al., 2007. Neurocognitive Effects of Methamphetamine: A Critical Review and Meta-analysis. *Neuropsychology Rev*, 17, pp.275–297.
- Sotak, C.H., 2002. The role of diffusion tensor imaging in the evaluation of ischemic brain injury - a review. *The role of diffusion tensor imaging in the evaluation of ischemic brain injury ± a review*, (15), pp.561–569.
- Tobias, M.C. et al., 2010. White-matter abnormalities in brain during early abstinence from methamphetamine abuse. *Psychopharmacology*, 209, pp.13–24.
- Wilson, J.M. et al., 1996. Striatal dopamine nerve terminal markers in human, chronic methamphetamine users. *Nature Medicine*, 2(6), pp.699–703.

## **CHAPTER 2: PUBLICATION-READY MANUSCRIPT**

# **WHITE MATTER CORRELATES OF NEUROPSYCHOLOGICAL FUNCTION IN ABSTINENT, YOUNG ADULT METHAMPHETAMINE USERS**

**Carla P Freeman<sup>a</sup>**

**Jonathan Ipser<sup>a</sup>**

**Jean-Paul Fouche<sup>b</sup>**

**Hetta Gouse<sup>a</sup>**

**Dan J Stein<sup>c</sup>**

**John A Joska<sup>a,d</sup>**

<sup>a</sup>Division of Neuropsychiatry, Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

<sup>b</sup>Department of Psychiatry, University of Stellenbosch, Stellenbosch, Western Cape, South Africa

<sup>c</sup>Head of Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

<sup>d</sup>Head of Clinical Services, Head of Division of Neuropsychiatry, Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

**Keywords:** methamphetamine; diffusion tensor imaging (DTI); fractional anisotropy; mean diffusion; white matter; neuropsychological function

## **Abstract**

### *Background*

Methamphetamine (MA) abuse is a global health concern due to widespread use and harmful effects, which includes neurotoxicity. This study aimed to describe neurocognitive deficits associated with MA dependence in young adults and to explore whether these deficits correlate with white matter (WM) microstructural abnormalities using diffusion tensor imaging (DTI).

### *Methods*

Twenty-one MA dependent individuals recently enrolled in an outpatient rehabilitation program and nineteen healthy controls participated in the study. Each participant completed a neuropsychological evaluation and underwent diffusion tensor imaging within one week of testing. Average whole-brain fractional anisotropy (FA) and mean diffusion (MD) measures derived from DTI data were compared between groups. Group differences in performance within specific neurocognitive domains and in a composite global neurocognitive score (GNS) were tested using non-parametric univariate statistics and within a linear regression framework, adjusting for age and gender. Correlation analyses were conducted to test associations between the neuropsychological data and selected frontal white matter (WM) tracts, including the genu and body of the corpus callosum (CC); right and left cingulum bundle (CB); right and left uncinate fasciculus (UF); right and left anterior corona radiata (CR) and the right and left superior longitudinal fasciculus (SLF).

### *Results*

No significant between-group differences were detected for performance in any of the neuropsychological domains assessed. No relationship between FA or MD and the GNS was demonstrated in the tracts of interest. After adjusting for age and gender, significant group differences in FA and MD were

detected across several regions of interest (ROI), however, these did not survive corrections for multiple comparisons.

### *Conclusion*

Cognitive performance and white matter integrity did not differ between young MA dependent subjects and healthy controls. Whatever differences were found in white matter did not survive correction for multiple comparisons. These findings may reflect one or more of several possibilities: that brain function and structure is relatively preserved in younger individuals; or that differences were too small to be detected in this sample. Further studies should explore the effects of aging, poly-substance abuse and HIV co-infection on neurocognitive functioning and structural brain integrity in methamphetamine users.

## **1. Introduction**

Methamphetamine (MA) is stimulant drug of abuse known to have detrimental effects on brain structure and function (Courtney & Ray 2014). MA presents a global health concern due to its addictive nature, widespread use and neurotoxic properties. Informal methamphetamine labs/kitchens are able to produce cheaper, more potent MA and continue to drive this problem (Maxwell & Brecht 2011; Brecht et al. 2004). South Africa, particularly the Western Cape province, is facing a growing MA epidemic (Plüddemann et al. 2013). According to the South African Community Epidemiology Network on Drug Use (SACENDU), 35% of individuals seeking treatment for drug use in 2006 report MA as their primary drug of choice (Plüddemann et al. 2008).

This highly lipophilic molecule dramatically enhances the release of catecholamines (dopamine; noradrenaline and serotonin) in the central nervous system and partially blocks their re-uptake. The increase in available catecholamines further stimulates the corresponding postsynaptic receptors (Cruickshank & Dyer 2009). Disruption of these neurotransmitter systems is thought to underlie the predominantly frontostriatal pattern of neurocognitive dysfunction previously reported in MA dependent individuals (Sekine et al.



2003). Deficits in episodic memory; executive function; motor skills; information processing speed and language have all been described in acute and chronic MA use (Maxwell & Brecht 2011; Scott et al. 2007). Some of these data suggest that decision-making may be impaired in MA dependent individuals, leading to high-risk behaviours and poor treatment outcomes (Pluddemann et al. 2008; Paulus et al. 2002). Intact neurocognitive function is important to treatment planning (many of which are cognitively-based interventions) and long-term rehabilitation.

This study aimed to investigate whether a young adult, MA dependent, South African population display neurocognitive deficits, and associated WM abnormalities detected on diffusion tensor imaging (DTI). DTI is a magnetic resonance imaging (MRI) modality, which allows for the non-invasive, in-vivo study of brain white matter (WM) structure based on the free diffusion of water molecules within WM tracts (Assaf & Pasternak 2007). The extent to which water molecules move within the same orientation within white matter fiber tracts, known as anisotropic diffusion, can be used to infer tissue microstructural abnormalities in certain disease states i.e. demyelination; inflammation; oedema; neoplasia and stroke (Alexander et al. 2007; Le Bihan et al. 2001).

Fractional anisotropy (FA) is a parameter of DTI and provides information about the orientation of white matter fibers within a voxel. FA is high (close to 1) if diffusivity is rapid and uninterrupted along fibers whereas FA approaches zero if diffusion occurs in all directions (Assaf & Pasternak 2007). Mean diffusivity (MD) provides an overall measure of diffusivity within a voxel or region and is understood to increase in pathological disease states (Le Bihan et al. 2001). For instance, in the MA DTI literature, Chung and colleagues demonstrated disruption of frontal structures in abstinent MA users in the anterior-posterior commissure (AC-PC) plane, which correlated with an impaired performance on the Wisconsin Card Sorting Test (WCST), a measure of executive functioning (Chung et al. 2007). Their group reported that male MA dependent subjects had significantly lower FA values in frontal white matter, compared to female MA participants and more errors on the

WCST. Kim et al. (2009) further demonstrated abnormal WM in the genu of the corpus callosum (CC) with corresponding impairment on the WCST. Lederer and colleagues demonstrated increased MD in the genu of the CC and right cingulum in chronic MA users (Lederer et al. 2015). DTI, combined with cognitive measures, could potentially provide a useful marker of MA induced alterations in human behaviour and response, for example, predicting treatment outcomes in a rehabilitation setting.

Our hypothesis was that WM abnormalities would follow a predominantly fronto-subcortical pattern in MA dependent individuals and would be associated with neurocognitive dysfunction represented by WM tracts in these areas. We predicted that group differences would be present in the following major frontal WM pathways: genu and body of the corpus callosum (CC) and right and left cingulum bundle (CB); right and left uncinate fasciculus (UF); the superior longitudinal fasciculus (SLF) and the anterior corona radiata (CR). The selection of the aforementioned WM tracts is based on the ICBM-DTI-81 white-matter atlas and represents tracts previously reported on in a separate cohort of methamphetamine patients by our lab (Mori et al. 2008; Uhlmann et al. 2016).

This study was approved by the University of Cape Town Human Research Ethics Committee (HREC no: 635/2014), and was conducted in accordance with the principles laid down by the Declaration of Helsinki 2013.

## **2. Methods and materials**

### **2.1. Subjects**

All MA dependent individuals, aged 18-40 years, attending the Tafelsig Matrix Rehabilitation Centre in Cape Town, South Africa, during the period of December 2011 to December 2012 were invited to participate in the study. Healthy, non-using, control subjects of a similar age were recruited in a convenience fashion from friends and family of MA + users living within the same community. Inclusion criteria for MA + participants were as follows: (1) completed 7 years of formal schooling; (2) right-handed; (3) able to converse

in English or Afrikaans and (5) MA dependence, as confirmed by the Mini International Neuropsychiatric Interview (MINI-500)(Sheehan et al. 1997).

Participants were excluded from the study if they were found to have the following: (1) a history of past or current serious mental illness (e.g. schizophrenia, bipolar mood disorder); (2) a history of past or current neurological illness affecting the brain (e.g. cerebrovascular illness, epilepsy); (3) severe systemic illness or significant head injury (loss of consciousness > 30mins and/or requiring an admission to hospital); (4) past or current substance abuse other than MA; alcohol; cannabis or nicotine; (5) HIV seropositivity; (6) contraindication to MRI and (7) refusal to sign informed consent. MA+ participants were required to abstain from MA use for 3 days prior to the neuropsychological assessment and 7 days prior to the neuroimaging. This was confirmed on self-report and by means of a urine drug screen (UDS) for amphetamines; opiates; cannabis and benzodiazepines. The Drug Use Disorders Identification Test (DUDIT), a sensitive, 11-item self-report tool, was used to screen for any substance use, in addition to methamphetamine (A. H. Berman et al. 2005). Healthy controls were excluded for any substance use, including cannabis. The DUDIT was used in conjunction with the Alcohol Use Disorders Identification Test (AUDIT), a widely used, validated 10-item tool allowing investigators to screen for excessive alcohol use (Bohn et al. 1995).

Measurement of total MA dose is notoriously difficult. In South Africa, MA is distributed in the form of straws at a cost of 40-60 ZAR per straw (the length of a drinking straw is filled with MA and divided into segments). Ten straws are estimated to hold an equivalent of 1 gram of MA, although this measurement may be inaccurate as the MA is often combined with other substances, such as talcum powder. For this study participants were asked to report dose as either number of straws or grams used per day or both. All participants signed full informed consent prior to entry in the study.

## 2.2. Neuropsychological assessment

The cognitive evaluation occurred on a single test day and was undertaken in the participant's first language (English or Afrikaans). The neuropsychological (NP) test battery was selected by a trained local neuropsychologist (HG) with expertise in MA abuse. The domains chosen for this study represent those most frequently associated with deficits in MA abuse (Scott et al. 2007; Gonzalez et al. 2004), and include the following tests, as listed in Table 2: (1) motor speed (Grooved Pegboard Test); (2) episodic memory (Hopkins Verbal Learning Test –Revised; total and delayed); (3) attention/working memory (WAIS – III Digit Span Subtest); (4) speed of information processing (Color Trails; WAIS – III Digit Symbol Subtest; Stroop color/word); (5) verbal fluency (Letter Fluency; Category Fluency) and (6) executive function (Stroop Color and Word; Color trails -II; Hayling Test).

### 2.3. Diffusion tensor imaging acquisition and processing

Magnetic resonance imaging was performed on a Siemens Magnetom 3T Allegra at the Cape Universities Brain Imaging Centre (CUBIC), within one week of the neuromedical and neuropsychological assessment. DTI data was acquired with the following parameters: TR=8800ms, TE=88ms, 30 diffusion-weighted volumes with  $b=1000\text{sec/mm}^2$  and 3 unweighted volumes ( $b=0\text{sec/mm}^2$ ). In-plane resolution was  $2 \times 2\text{ mm}^2$  with a slice thickness of 2.2 mm. The sequence was repeated 3 times to improve signal-to-noise ratio. In order to facilitate registration of the DTI data in standard space, thereby allowing group comparisons of DTI outcomes, a high resolution multi-echo MPRAGE T1-weighted image was acquired, with the following parameters: TR=2530ms; TE=1.64ms, 3.5ms, 5.36ms and 7.22ms; FOV=256x256; 176 slices, 1mm isotropic voxel size.

The 3  $b=0\text{sec/mm}^2$  images for each DTI repetition were averaged to improve the signal-to-noise ratio. Eddy current correction was performed with the FMRIB Software Library (FSL 4.1.8) on each acquisition (Smith et al. 2006). Data were exported to Matlab (Mathworks Inc, Natick, MA) for further processing with all three acquisitions being co-registered by using the  $b=0\text{ mm/s}^2$  as reference. The DTI acquisitions were averaged and exported to FSL for tensor fitting and brain extraction. Outliers were calculated by obtaining the

Z-value of the tensor estimates at the 25<sup>th</sup> and 75<sup>th</sup> percentile. Outliers were discarded if the data points were more than 3 standard deviations from the mean. Whole brain statistical analysis was undertaken using FSL's tract-based spatial statistics (TBSS) software program (Smith et al. 2006). The surface data produced by Freesurfer was smoothed and imported into Freesurfer's qdec toolbox for vertex-level analysis. The following ROIs were selected based on a predefined mask as per the ICBM-DTI-18 white matter-labels atlas: genu and body of the corpus callosum (CC) and bilaterally for the cingulum bundle (CB); the uncinate fasciculus (UF); the superior longitudinal fasciculus (SLF) and the corona radiata (CR)(Mori et al. 2008).

#### 2.4. Statistical analysis

MA + users were compared with healthy controls with respect to demographic and clinical characteristics. The effect of gender on these sample characteristics was also determined. Group differences on categorical data were assessed using chi-square tests. Continuous outcomes were analysed using independent t-tests, or alternatively, non-parametric two-sided Mann Whitney or Kruskal-Wallis tests in cases in which parametric assumptions of normality and heteroscedascity were not confirmed.

Neurocognitive domains scores were calculated by normalising the raw scores from constituent tasks by means of their conversion into Z scores, and then averaging the resulting scores across the tasks within each domain. The rationale behind creating Z scores is to have standardized scores on the same scale which can be compared across domains. Z scores were obtained by subtracting the mean of the scores of the healthy control group from the mean of the MA + group and dividing the result by the standard deviation of the control group. i.e.  $X_{MA+} - X_{MA-} / SD(MA-)$ . In addition, a global neurocognitive score (GNS) was derived from the mean Z score across all of the domains for each participant. Group differences in domain scores were determined using independent samples t-tests. In order to adjust for the effects of age and gender on domain scores, test estimates for between-group comparisons on the domain scores were also extracted from linear regression models with group membership, age and gender as predictor variables. In addition, non-

parametric spearman's correlations were calculated between the FA and MD measures from the tracts of interest and the GNS within each of the groups. The risk of false positives was minimised by employing bonferroni correction for multiple comparisons

All statistical analyses were conducted using the R statistical computing platform (version 3.2; R Core Team) with alpha set to 0.05 (RCoreTeam 2013).

### **3. Results**

#### **3.1. Demographic data**

21 MA abusers (48% male, mean age = 26.7 years) currently engaged in the MATRIX rehabilitation program and 19 healthy controls (26% male, average age = 27.4 years) participated in this research study. The study groups were comparable with respect to age, education, housing and relationship status (see Table 1). There was no difference in alcohol use between the two groups as measured on the AUDIT ( $p>0.1$ ). There was preliminary evidence that concurrent cannabis use was greater in male MA subjects, though this was not statistically significant ( $p=0.06$ ).

#### **3.2 MA use characteristics**

The average age of first MA use was 19.6 years of age. Most MA users engaged in daily use of the drug. Males and females did not differ with respect to age of first use of MA, duration of use or abstinence, amount of money spent on MA per week, and the frequency or estimated amount of MA used (see Table 1).

#### **3.3. Neuropsychological testing**

No statistically significant differences in neurocognitive performance were found between MA dependent individuals and healthy comparison subjects for any of the cognitive domains (Table 2).

### 3.4. Diffusion tensor imaging analysis

#### 3.4.1 Effect of MA on mean diffusion (MD)

Regression models of group differences for the frontal tracts of particular interest in this study (genu and body of the CC, bilateral cingulum bundle; bilateral uncinate fasciculus; bilateral anterior corona radiata; bilateral superior longitudinal fasciculus) revealed a sub-threshold increase in MD in the right uncinate fasciculus ( $t = -1.79$ ,  $p = 0.08$ ), after adjusting for the effects of age and gender (Table 3). Age had a significant impact on several ROIs after adjusting for gender and MA use. These, however, did not survive bonferroni correction for multiple comparisons.

Given lack of evidence of group differences in the planned comparisons, post-hoc exploratory univariate comparisons of mean diffusion for forty-eight ROIs between the groups was conducted. These additional analyses detected significantly increased MD in the external capsule only (unadjusted  $p = 0.02$ ) in MA participants (Table S2).

#### 3.4.2 Effect of MA on fractional anisotropy (FA)

There was no evidence from regression models for group differences between the tracts of interest (Table 3). Sub-threshold evidence of a decrease of FA in the left superior longitudinal fasciculus relative to the control group was observed in MA+ patients, however ( $t = 1.96$ ,  $p = 0.06$ ).

Additional post-hoc exploratory univariate DTI comparisons for the 48 ROIs revealed no significant differences in FA between MA dependent users and healthy controls (Table S1). Greater age was associated with a significant reduction in FA in the right uncinate fasciculus ( $p=0.04$ ), after adjusting for gender and group status.

#### 3.4.3 Relationship between neuropsychological performance and DTI in selected ROIs

Greater FA in the left uncinate fasciculus was associated with an impaired global neurocognitive score in MA dependent individuals ( $Rho = -0.48$ ,  $p = 0.04$ ), but not in controls ( $Rho = 0.05$ ,  $p = 0.85$ ; see Table 4), a difference which did not survive correction for multiple comparisons. MD scores were not informative with respect to cognitive impairment for any of the ROIs.

#### **4. Discussion**

In this pilot study of young adult, MA dependent users, we were not able to demonstrate significant differences in neuropsychological performance or white matter abnormalities on DTI when compared with healthy control participants. No significant between-group findings for both FA and MD were observed in our tracts of interest. After adjusting for age and gender, though, a sub-threshold decreased FA was observed in the left superior longitudinal fasciculus in MA+ patients, and increased MD in the right uncinate fasciculus. The above findings were consistent with our hypothesis and in keeping with reported DTI findings in the MA literature. In addition, increased FA in the left uncinate fasciculus was associated with poorer neuropsychological performance, as measured by a global neurocognitive score, however, none of these findings survived correction for multiple comparisons.

A number of factors may have contributed to the negative findings of this study. The average age of our study sample (26.7 years) was considerably younger than populations reported in the MA / DTI literature. For instance, Salo and colleagues (2009) demonstrated WM abnormalities in the CC, correlating with poor cognitive control, in MA users with an average age of 36.29 ( $\pm 8.7$ ) years. Kim et al. (2009) reported on MA dependent subjects with an average age of 34.4 ( $\pm 2.9$ ) years and showed decreased FA in the CC, which corresponded with a worse performance on the Wisconsin Card Sorting Test (WCST). Younger age may be protective in MA use, raising the possibility of an initial adaptive response to MA due to increased neuronal plasticity in younger brains, as suggested by Berman and colleagues (Berman et al. 2008). Furthermore, a temporary imbalance in the frontostriatal circuits due to differences in maturation rates between the cortex and striatum has



been suggested by Vink and colleagues (Vink et al. 2014). This may serve to explain some variability in the neuropsychological task performance. Our sample also had a relatively shorter duration of MA use in comparison to other study populations ( $7.4 \pm 3.65$  vs.  $10.6 \pm 2.7$  years in Kim et al. (2009)), which may have impacted on the severity of neurotoxicity and subsequent WM damage. Furthermore, the effects of gender should be considered, particularly with regards to MA use characteristics. Although our sample was small and differences between male and female participants were not significant, male MA users engaged in more frequent MA use; had a longer duration of use and were more likely to abuse cannabis. A larger sample may reflect both DTI and neuropsychological differences associated with these differences.

In keeping with Chung et al. we expected to find lower FA values in the prefrontal and frontal cortices (Chung et al. 2007). Microstructural WM abnormalities are present within 7-13 days of abstinence from MA, notably in the prefrontal cortex and hippocampal formation (Tobias et al. 2010). The duration of abstinence in our sample ranged from an average of 38.8 days (range: 10-65 days) in male subjects to 60 days (range: 24-126 days) in female MA users. The absence of WM abnormalities in this sample is unexpected and may indicate the possibility of neuronal recovery during the abstinent period. Nordahl et al. (2005) suggest a period of normalization of neuronal structure and function following MA abstinence, supported by the normalization of choline levels on magnetic resonance spectroscopy in the anterior cingulum. Wang et al demonstrated recovery of thalamic metabolism in abstinent MA users and suggest that this may reflect a compensatory response to the dopaminergic deficit present in the striatum, subsequently improving neuropsychological performance (Wang 2004). Furthermore Volkow et al. (2001) demonstrated the recovery of the dopamine transporter following protracted abstinence from MA. FA is a measure of the movement of water molecules within WM tracts. FA is thought to increase when this diffusion is highly ordered within intact WM. Our finding of a sub-threshold positive correlation between increased FA in the left superior fasciculus and a measure of global cognitive performance in healthy controls supports this interpretation of FA as an index of WM integrity. The cingulum forms part of

the frontal-striatal network of WM, essential for higher-order executive functioning. Nevertheless we were unable to demonstrate a significant deficit in executive function in MA dependent individuals.

### Limitations

The sample size of this study is small, though comparable to other DTI studies of MA use (Chung et al. 2007, Kim et al. 2009), and a larger sample may have provided sufficient power to yield positive findings. We were not able to include the WCST in our neuropsychological test battery, known to be associated with DTI abnormalities (Chung 2007, Kim 2009), due to time and resource constraints. MA users are expected to have comparatively lower mean scores for the neurocognitive tests, yet may still perform within the normed limits for their age and education (Hart et al. 2011). The lack of group differences in test performance may reflect a selection bias within the healthy comparison group. The healthy control group was selected predominantly from family and friends accompanying the MA dependent user to their rehab visits and alcohol use was not excluded. Female healthy controls ( $n = 14$ ) were mostly unemployed and had the highest scores on the AUDIT, which may, in turn, have contributed to impaired test scores.

Due to the frequent co-morbidity of co-morbid cannabis use in this population, we elected not to exclude cannabis users for the MA+ group. Male MA users trended towards an increased co-morbid use of cannabis ( $p=0.06$ ). Although not statistically significant, a larger sample size may have highlighted this potential difference between genders with implications for between group differences. Gonzalez et al. (2004) postulated a protective effect of cannabis on neurocognitive performance in MA abuse, yet were unable to demonstrate this. The authors concluded that comorbid cannabis use did not exacerbate the neurotoxic properties of MA, and recommended further studies of the potentially neuro-protective qualities of cannabis in the context of MA use (Gonzalez et al. 2004).

Dosage is an important aspect of understanding the neurotoxic potential of MA. We attempted to determine an estimate of the dosage (based on self-

report by MA dependent individuals), however, we feel that these estimates are highly inaccurate as MA in South Africa is frequently cut and sold with other inert substances and is therefore difficult to measure. Although the authors recommend a more accurate measurement of MA potency in future studies, we are aware that this is a near impossible task due to the variability of substance produced in informal MA labs.

## Conclusion

The absence of neuropsychological or white matter deficits in this relatively young adults group of MA dependent individuals may suggest a neuroprotective aspect associated with younger age. There may indeed be a tipping-point, whereby neurotoxicity is reflected on cognitive measures and neuroimaging. The authors recommend further research examining the effects of age; co-morbid substance abuse and HIV co-infection using a matched-control group.

## Acknowledgements

This work was supported, in part, by a grant from the Discovery Foundation. Dr. Freeman is a 2010 recipient of a Discovery Foundation Academic Award.

## Declaration of conflicts of interest

The authors report no conflicts of interest.

## References

- Alexander, A. et al., 2007. Diffusion tensor imaging of the brain. *Neurotherapeutics*, 4(3), pp.316–329.
- Assaf, Y. & Pasternak, O., 2007. Diffusion Tensor Imaging (DTI)-based White Matter Mapping in Brain Research: A Review. *J Mol Neurosci*, 34, pp.51–61.
- Berman, A.H. et al., 2005. Evaluation of the Drug Use Disorders Identification Test (DUDIT) in Criminal Justice and Detoxification Settings and in a Swedish Population Sample. *European Addiction Research*, 11(1), pp.22–31.
- Berman, S. et al., 2008. Abuse of Amphetamines and Structural Abnormalities in the Brain. *Annals of the New York Academy of Sciences*, 1141(1),

pp.195–220.

- Bohn, M.J., Babor, T.F. & Kranzler, H.R., 1995. The alcohol use disorders identification test (AUDIT): Validation of a screening instrument for use in medical settings\*. *Journal of Studies on Alcohol*, pp.423–432.
- Brecht, M.L. et al., 2004. Methamphetamine use behaviors and gender differences. *Addictive Behaviors*, 29(1), pp.89–106.
- Chung, A. et al., 2007. Decreased frontal white-matter integrity in abstinent methamphetamine abusers. *International Journal of Neuropsychopharmacology*, 10, pp.765–775.
- Courtney, K.E. & Ray, L.A., 2014. Methamphetamine: An update on epidemiology, pharmacology, clinical phenomenology, and treatment literature. *Drug and Alcohol Dependence*, 143, pp.11–21.
- Cruickshank, C.C. & Dyer, K.R., 2009. A review of the clinical pharmacology of methamphetamine. *Addiction*, 104(7), pp.1085–1099.
- Gonzalez, R. et al., 2004. Neurocognitive performance of methamphetamine users discordant for history of marijuana exposure. *Drug and Alcohol Dependence*, 76(2), pp.181–190.
- Hart, C.L. et al., 2011. Is Cognitive Functioning Impaired in Methamphetamine Users? A Critical Review. *Neuropsychopharmacology*, pp.1–23.
- Kim, I.-S. et al., 2009. Reduced corpus callosum white matter microstructural integrity revealed by diffusion tensor eigenvalues in abstinent methamphetamine addicts. *Neurotoxicology*, 30, pp.209–213.
- Kirkpatrick, M.G. et al., 2007. Effects of intranasal methamphetamine on metacognition of agency. *Psychopharmacology*, 197(1), pp.137–144.
- Le Bihan, D. et al., 2001. Diffusion Tensor Imaging: Concepts and Applications. *Journal Of Magnetic Resonance Imaging*, 13, pp.534–546.
- Lederer, K. et al., 2015. Frontal white matter changes and aggression in methamphetamine dependence. *Metabolic Brain Disease*, 31(1), pp.53–62.
- Maxwell, J.C. & Brecht, M.-L., 2011. Methamphetamine: Here we go again? *Addictive Behaviors*, 36(12), pp.1168–1173.
- Mori, S. et al., 2008. Stereotaxic White Matter Atlas Based on Diffusion Tensor Imaging in an ICBM Template. *NeuroImage*, 40(2), pp.570–582.
- Nordahl, T.E. et al., 2005. Methamphetamine Users in Sustained Abstinence. *Arch Gen Psychiatry*, 62, pp.444–452.
- Paulus, M.P. et al., 2002. Behavioral and Functional Neuroimaging Evidence for Prefrontal Dysfunction in Methamphetamine-Dependent Subjects.

- Neuropsychopharmacology*, 26(1), pp.53–63.
- Pluddemann, A., Myers, B.J. & Parry, C., 2008. Surge in treatment admissions related to methamphetamine use in Cape Town, South Africa: implications for public health. *Drug and Alcohol Review*, 27, pp.185–189.
- Plüddemann, A. et al., 2013. Monitoring the prevalence of methamphetamine-related presentations at psychiatric hospitals in Cape Town, South Africa. *African Journal of Psychiatry*, 16(1).
- RCoreTeam, 2013. R: A language and environment for statistical computing. [www.R-project.org](http://www.R-project.org). Available at: <http://www.R-project.org/>.
- Salo, R. et al., 2009. Cognitive Control and White Matter Callosal Microstructure in Methamphetamine Dependent Subjects: A DTI Study. *Biological Psychiatry*, 65(2), pp.122–128.
- Scott, J.C. et al., 2007. Neurocognitive Effects of Methamphetamine: A Critical Review and Meta-analysis. *Neuropsychology Rev*, 17, pp.275–297.
- Sekine, Y. et al., 2003. Association of Dopamine Transporter Loss in the Orbitofrontal and Dorsolateral Prefrontal Cortices With Methamphetamine-Related Psychiatric Symptoms. *Am J Psychiatry*, 160, pp.1699–1701.
- Sheehan, D.V. et al., 1997. Reliability and Validity of the M.I.N.I. International Neuropsychiatric Interview (M.I.N.I.): According to the SCID-P. *European Psychiatry*, 12, pp.232–241.
- Smith, S.M. et al., 2006. Tract-based spatial statistics: Voxelwise analysis of multi-subject diffusion data. *NeuroImage*, 31, pp.1487–1505.
- Tobias, M.C. et al., 2010. White-matter abnormalities in brain during early abstinence from methamphetamine abuse. *Psychopharmacology*, 209, pp.13–24.
- Uhlmann, A. et al., 2016. White matter microstructure and impulsivity in methamphetamine dependence with and without a history of psychosis. *Hum Brain Mapp*, pp.00:00-00.
- Vink, M. et al., 2014. Frontostriatal activity and connectivity increase during proactive inhibition across adolescence and early adulthood. *Hum Brain Mapp*, 12, pp.4415–4427.
- Volkow, N.D. et al., 2001. Association of Dopamine Transporter Reduction With Psychomotor Impairment in Methamphetamine Abusers. *Am J Psychiatry*, 158, pp.377–382.
- Wang, G. et al., 2004. Partial Recovery of Brain Metabolism in Methamphetamine Abusers after Protracted Abstinence. *Am J Psychiatry*, 161: pp.242-248.

**Table 1: Participant demographic data and clinical characteristics**

Description	MA+ (n=21)		MA - (n=19)		Test statistics <sup>a</sup>
	Male (n=10)	Female (n=11)	Male (n=5)	Female (n=14)	
Age, years, mean (SD)	28.55 (5.43)	25.05 (4.70)	29.59 (5.97)	26.66 (5.32)	t = 2.91,(3), p = 0.41
Years of formal education, mean (SD)	10.2 (2.49)	10.27 (2.2)	10.2 (2.49)	10.21 (1.31)	t = 0.23,(3), p = 0.97
Unemployed, %	10 (100%)	9 (82%)	2 (40%)	10 (71%)	$\chi^2 = 8.89, (4),$ p = 0.06
Single, %	9 (90%)	9 (82%)	3 (60%)	7 (50%)	$\chi^2 = 8.27, (4),$ p = 0.08
Living in family home, %	9 (90%)	9 (82%)	2(40%)	8 (57%)	$\chi^2 = 7.77, (4),$ p = 0.10
Audit mean, (SD)	1.7, (2.1)	0.8, (1.7)	0.6, (1.3)	2.4,(3.1)	t = 3.18, (3), p = 0.36
Dudit mean, (SD)	21.4, (8.9)	19.3, (8.6)	0 (0)	0 (0)	z = -1.10, (1), p = 0.27
<b>MA use characteristics</b>					
Age, in years, of first use, mean, (SD)	20, (5.54)	19.11, (4.83)	-	-	Z=-1.10, p = 0.71
Weekly cost of MA, %: <R500 R500 to R1000 R1000 to R2000 >R2000	5 (50%) 2 (20%) 2 (20%) 1 (10%)	8 (89%) 1 (11%) - -	-	-	$\chi^2 = 3.98, (3),$ p = 0.26
Frequency of use Daily, % Weekly, %	9 (90%) 1 (10%)	6 (67%) 3 (33%)	-	-	$\chi^2 = 1.55, (1),$ p = 0.21
Daily dose in grams, mean, (SD)	2.17, (2.11)	1.51, (1.83)	-	-	Z = -1.07, p = 0.28
Duration of use, yrs, mean, (SD)	8.01, (4.13)	6.72, (3.12)	-	-	Z = -0.53, p = 0.59
Days abstinent, mean, (SD)	38.8, (23.0)	60.9, (38.9)	-	-	Z = 1.09, p = 0.27
Co-morbid cannabis use, %	80	40	-	-	$\chi^2 = 3.33, (1),$ p = 0.06

Note: MA = methamphetamine; SD = standard deviation; AUDIT = Alcohol Use Disorders Identification Test; DUDIT = Drug Use Disorders Identification Test; IQR = interquartile range  
<sup>a</sup> degrees of freedom in parentheses

**Table 2: Neuropsychological performance of MA users and healthy comparison subjects**

Variable <sup>a</sup>	MA + (n = 21)	MA – (n = 19)	Statistic	Adjusted statistic <sup>b</sup>
<b>Motor speed</b>				
GP Dominant	68.1 (7.87)	69.32 (8.49)		
GP Non-dominant	71.86 (9.06)	71.26 (8.69)		
Motor speed Z	0.02 (0.89)	-0.02 (0.87)	t = -0.15 (37.78), p = 0.88	t = 1.30, p = 0.20
<b>Episodic memory</b>				
HVLT-R total	20.86 (4.08)	20.89 (4.25)		
HVLT-R delay	7.29 (2)	6.47 (2.55)		
Episodic memory Z	0.08 (0.87)	-0.09 (1.01)	t = -0.58 (35.75), p = 0.57	t = 0.64, p = 0.53
<b>Attention/working memory</b>				
Digit span	11.29 (3.3)	12.11 (2.81)		
Working memory Z	-0.13 (1.0)	0.14 (0.92)	t = 0.85 (37.9), p = .40	t = -1.01, p = 0.32
<b>Speed of information processing</b>				
Colour trails I	51 (16.64)	43.26 (9.8)		
Digit symbol	45.48 (11.49)	50.58 (12.78)		
Stroop Color Naming	45.71 (14.54)	38.16 (11.62)		
Info speed Z	-0.12 (0.85)	0.14 0.49	t = 1.22 (32.51), p = 0.23	t = -1.07. p = 0.29
<b>Verbal fluency</b>				
COWAT – FAS	0.17 (0.99)	-0.19 (0.79)		
Animal fluency	33.76 (7.83)	32.37 (7.15)		
Fluency Z	0.17 (0.99)	-0.19 (0.79)	t = -1.25 (37.44), p = 0.22	t = 1.24, p = 0.22
<b>Executive function <sup>c</sup></b>				
Stroop Color and Word Test	45.71 (14.54)	38.16 (11.62)		
Colour trails 2	92.76 (26.05)	88.11 (18.61)		
Hayling Test	6.4 (3.68)	7.47 (4.59)		
Executive function Z	0.03 (0.77)	-0.02 (0.68)	t = -0.19 (36.80), p = 0.85	t = 0.13, p = 0.90

Note: <sup>a</sup>All scores represent raw scores, presented as means and standard deviations. Higher domain Z scores reflect better test performance. <sup>b</sup> Group comparisons after adjusting for the effects of gender and age within a linear regression model <sup>c</sup> Data was missing for one MA + participant.

MA + = Methamphetamine user; MA - = healthy comparison subject

HVLT-R = Hopkins Verbal Learning Test – Revised; COWAT – Controlled Oral Word Association Test; GP – Grooved Pegboard

**Table 3: Differences between MA+ and MA- groups on DTI outcomes for tracts of interest**

Region of interest	MA+ (M $\pm$ SD)	MA- (M $\pm$ SD)	Model est. $\pm$ SD	T-test statistic	alpha
<b>FA</b>					
Genu CC	0.39 $\pm$ 0.04	0.40 $\pm$ 0.03	0,014 $\pm$ -0,012	t = 1,21	p = 0,23
Body CC	0.39 $\pm$ 0.04	0.39 $\pm$ 0.02	0,006 $\pm$ -0,011	t = 0,48	p = 0,63
R Cingulum	0.38 $\pm$ 0.04	0.40 $\pm$ 0.03	0,020 $\pm$ -0,012	t = 1,61	p = 0,12
L Cingulum	0.41 $\pm$ 0.03	0.41 $\pm$ 0.02	0,004 $\pm$ -0,009	t = 0,49	p = 0,63
R Uncinate fasciculus	0.32 $\pm$ 0.05	0.33 $\pm$ 0.04	0,006 $\pm$ -0,015	t = 0,37	p = 0,71
L Uncinate fasciculus	0.45 $\pm$ 0.04	0.45 $\pm$ 0.02	0,007 $\pm$ -0,010	t = 0,66	p = 0,51
R Ant corona radiata	0.49 $\pm$ 0.04	0.49 $\pm$ 0.03	0,009 $\pm$ -0,012	t = 0,78	p = 0,44
L Ant corona radiata	0.48 $\pm$ 0.04	0.49 $\pm$ 0.02	0,005 $\pm$ -0,010	t = 0,53	p = 0,60
R Sup long fasciculus	0.38 $\pm$ 0.04	0.37 $\pm$ 0.04	-0,009 $\pm$ -0,013	t = -0,73	p = 0,47
L Sup long fasciculus	0.34 $\pm$ 0.04	0.36 $\pm$ 0.02	0,019 $\pm$ -0,010	t = 1,96	p = 0,06*
<b>MD</b>					
Genu CC	0.75 $\pm$ 0.03	0.75 $\pm$ 0.03	0,051 $\pm$ -0,103	t = 0,49	p = 0,62
Body CC	0.72 $\pm$ 0.04	0.73 $\pm$ 0.02	0,056 $\pm$ -0,0965	t = 0,58	p = 0,56
R Cingulum	0.82 $\pm$ 0.03	0.82 $\pm$ 0.03	-0,048 $\pm$ -0,091	t = -0,53	p = 0,60
L Cingulum	0.70 $\pm$ 0.03	0.70 $\pm$ 0.03	-0,057 $\pm$ -0,082	t = -0,69	p = 0,49
R Uncinate fasciculus	1.38 $\pm$ 0.15	1.31 $\pm$ 0.13	-0,759 $\pm$ -0,425	t = -1,79	p = 0,08*
L Uncinate fasciculus	0.68 $\pm$ 0.03	0.68 $\pm$ 0.03	-0,027 $\pm$ -0,105	t = -0,26	p = 0,79
R Ant corona radiata	0.93 $\pm$ 0.08	0.94 $\pm$ 0.06	0,031 $\pm$ -0,238	t = 0,13	p = 0,89
L Ant corona radiata	0.83 $\pm$ 0.06	0.82 $\pm$ 0.04	-0,058 $\pm$ -0,157	t = -0,37	p = 0,71
R Sup long fasciculus	0.81 $\pm$ 0.08	0.83 $\pm$ 0.09	0,368 $\pm$ -0,268	t = 1,37	p = 0,18
L Sup long fasciculus	1.48 $\pm$ 0.14	1.43 $\pm$ 0.13	-0,630 $\pm$ -0,435	t = -1,45	p = 0,16

Model coefficients extracted from multiple linear regression of DTI outcomes on group membership, adjusting for differences in gender and age.

MD values multiplied by 1000

\*p<0.1, \*\*p<0.05



**Table 4: DTI correlates of Global Neurocognitive Score**

Region of interest	rho MA+	alpha	rho MA-	alpha
<b>FA</b>				
Genu CC	-0.356	p = 0.12	-0.03	p = 0.92
Body CC	-0.30	p = 0.19	0.04	P = 0.88
R Cingulum	-0.14	p = 0.56	0.31	p = 0.20
L Cingulum	-0.26	p = 0.28	-0.08	p = 0.75
R Uncinate fasciculus	-0.17	p = 0.48	-0.14	p = 0.58
L Uncinate fasciculus	-0.48	p = 0.04	0.05	p = 0.85
R Anterior Corona Radiata	-0.15	p = 0.52	0.36	p = 0.13
L Anterior Corona Radiata	-0.24	p = 0.31	-0.04	p = 0.87
R Superior longitudinal fasciculus	-0.23	p = 0.33	0.20	p = 0.42
L Superior longitudinal fasciculus	-0.29	p = 0.21	0.05	p = 0.85
<b>MD</b>				
Genu CC	0.32	p = 0.17	0.05	p = 0.83
Body CC	0.29	p = 0.22	-0.12	p = 0.62
R Cingulum	0.21	p = 0.38	-0.03	p = 0.91
L Cingulum	0	p = 1	-0.07	p = 0.79
R Uncinate fasciculus	0.09	p = 0.72	0.36	p = 0.13
L Uncinate fasciculus	-0.7	p = 0.78	0.03	p = 0.90
R Anterior Corona Radiata	0.38	p = 0.10	-0.11	p = 0.66
L Anterior Corona Radiata	0.26	p = 0.26	0.07	p = 0.78
R Superior longitudinal fasciculus	0.14	p = 0.55	-0.45	p = 0.05
L Superior longitudinal fasciculus	0.22	p = 0.35	0.01	p = 0.97

MA+ = MA dependent

Bonferroni corrected p = 0.00

## Supplementary material

**Table S1: Mean DTI values in MA dependent individuals and healthy control subjects**

Region	FA Ma +	FA MA-	FA: t value, p value	MD* MA+	MD* MA-	MD: t value, p value
Middle cerebellar peduncle	0.43 ±0.04	0.43 ±0.02	t = -0.75, p = 0.46	0.71 ±0.03	0.71 ±0.03	t = -0.08, p = 0.93
Pontine crossing tract	0.47 ±0.04	0.47 ±0.02	t = -0.64, p = 0.54	0.69 ±0.02	0.68 ±0.02	t = 1.08, p = 0.29
Genu CC	0.39 ±0.04	0.40 ±0.03	t = -1.12, p = 0.27	0.75 ±0.03	0.75 ±0.03	t = -0.23, p = 0.82
Body CC	0.39 ±0.04	0.39 ±0.02	t = -0.29, p = 0.78	0.72 ±0.04	0.73 ±0.02	t = -0.41, p = 0.68
Splenium CC	0.45 ±0.03	0.45 ±0.03	t = -0.04, p = 0.97	0.96 ±0.07	0.96 ±0.03	t = 0.08, p = 0.94
Fornix	0.43 ±0.03	0.43 ±0.02	t = -0.36, p = 0.72	1.03 ±0.08	1.03 ±0.04	t = 0.09, p = 0.93
R Corticospinal tract	0.56 ±0.04	0.56 ±0.02	t = -0.41, p = 0.69	0.75 ±0.03	0.74 ±0.02	t = 1.09, p = 0.28
L Corticospinal tract	0.55 ±0.04	0.55 ±0.02	t = -0.52, p = 0.61	0.73 ±0.02	0.72 ±0.02	t = 0.63, p = 0.53
R Medial lemniscus	0.47 ±0.04	0.47 ±0.03	t = -0.14, p = 0.89	0.66 ±0.04	0.66 ±0.03	t = 0.15, p = 0.88
L Medial lemniscus	0.46 ±0.03	0.46 ±0.02	t = -0.60, p = 0.55	0.67 ±0.03	0.68 ±0.02	t = -0.74, p = 0.47
R Inf cerebellar peduncle	0.54 ±0.03	0.54 ±0.02	t = 0.02, p = 0.99	0.69 ±0.03	0.69 ±0.02	t = 0.26, p = 0.79
L Inf cerebellar peduncle	0.36 ±0.04	0.37 ±0.02	t = -1.03, p = 0.31	0.67 ±0.03	0.67 ±0.03	t = 0.36, p = 0.72
R Sup cerebellar peduncle	0.54 ±0.03	0.54 ±0.01	t = -0.28, p = 0.78	0.70 ±0.02	0.70 ±0.02	t = -1.18, p = 0.25
L Sup cerebellar peduncle	0.49 ±0.04	0.49 ±0.03	t = 0.22, p = 0.82	0.74 ±0.03	0.74 ±0.02	t = 0.39, p = 0.71
R Cerebral peduncle	0.47 ±0.04	0.47 ±0.03	t = -0.65, p = 0.52	0.74 ±0.03	0.74 ±0.03	t = 0.06, p = 0.95
L Cerebral peduncle	0.39 ±0.03	0.39 ±0.03	t = 0.08, p = 0.94	0.72 ±0.03	0.72 ±0.03	t = -0.74, p = 0.47
R Ant limb of int capsule	0.39 ±0.03	0.39 ±0.02	t = 0.19, p = 0.84	0.72 ±0.03	0.73 ±0.03	t = -0.72, p = 0.48
L Ant limb of int capsule	0.42 ±0.03	0.43 ±0.02	t = -0.89, p = 0.38	0.69 ±0.03	0.69 ±0.02	t = 0.28, p = 0.78
R Post limb of int capsule	0.41 ±0.03	0.41 ±0.02	t = -0.25, p = 0.80	0.69 ±0.03	0.73 ±0.03	t = -1.11, p = 0.28
L Post limb of int capsule	0.42 ±0.03	0.42 ±0.03	t = -0.79, p = 0.44	0.78 ±0.04	0.78 ±0.04	t = 0.14, p = 0.89
R Retrolenticular int capsule	0.39 ±0.03	0.40 ±0.02	t = -0.72, p = 0.48	0.79 ±0.04	0.78 ±0.03	t = 0.32, p = 0.75
L Retrolenticular int capsule	0.51 ±0.04	0.51 ±0.02	t = 0.25, p = 0.81	0.81 ±0.06	0.80 ±0.04	t = 1.09, p = 0.28
R Ant corona radiata	0.49 ±0.04	0.49 ±0.03	t = -0.48, p = 0.63	0.93 ±0.08	0.94 ±0.06	t = -0.12, p = 0.90
L Ant corona radiata	0.48 ±0.04	0.49 ±0.02	t = -0.45, p = 0.65	0.83 ±0.06	0.82 ±0.04	t = 0.28, p = 0.78
R Sup corona radiata	0.46 ±0.03	0.45 ±0.03	t = 0.45, p = 0.65	0.83 ±0.04	0.82 ±0.03	t = 1.07, p = 0.29
L Sup corona radiata	0.41 ±0.03	0.41 ±0.02	t = 0.12, p = 0.90	0.81 ±0.04	0.81 ±0.04	t = 0.70, p = 0.48
R Post corona radiata	0.34 ±0.03	0.34 ±0.02	t = 0.01, p = 0.99	0.73 ±0.03	0.72 ±0.02	t = 0.09, p = 0.92
L Post corona radiata	0.33 ±0.03	0.33 ±0.02	t = -0.23, p = 0.82	0.75 ±0.03	0.76 ±0.02	t = -1.44, p = 0.15
R Post thalamic radiation	0.40 ±0.04	0.39 ±0.03	t = 0.57, p = 0.57	0.72 ±0.03	0.72 ±0.02	t = 0.73, p = 0.47
L Post thalamic radiation	0.42 ±0.04	0.41 ±0.03	t = 0.22, p = 0.83	0.70 ±0.03	0.70 ±0.02	t = -0.17, p = 0.87
R Sagittal stratum	0.29 ±0.05	0.29 ±0.04	t = -0.02, p = 0.98	0.81 ±0.04	0.81 ±0.05	t = -0.19, p = 0.84
L Sagittal stratum	0.28 ±0.05	0.29 ±0.03	t = -0.31, p = 0.76	0.82 ±0.06	0.82 ±0.05	t = 0.47, p = 0.64
R Ext capsule	0.38 ±0.04	0.38 ±0.04	t = -0.44, p = 0.66	0.91 ±0.03	0.89 ±0.03	t = 2.33, p = 0.02*
L Ext capsule	0.50 ±0.07	0.53 ±0.03	t = -1.44, p = 0.16	0.88 ±0.07	0.86 ±0.03	t = 1.30, p = 0.20
R Cingulum	0.38 ±0.04	0.40 ±0.03	t = -1.49, p = 0.14	0.82 ±0.03	0.82 ±0.03	t = 0.29, p = 0.77
L Cingulum	0.41 ±0.03	0.41 ±0.02	t = -0.21, p = 0.84	0.70 ±0.03	0.70 ±0.03	t = 0.25, p = 0.80
R Cingulum hippocampus	0.38 ±0.02	0.39 ±0.02	t = -0.85, p = 0.40	0.71 ±0.02	0.72 ±0.02	t = -0.86, p = 0.39
L Cingulum hippocampus	0.40 ±0.05	0.41 ±0.03	t = -0.72, p = 0.47	0.66 ±0.03	0.65 ±0.02	t = 0.61, p = 0.54
R Fornix stria terminalis	0.38 ±0.05	0.38 ±0.03	t = -0.15, p = 0.88	0.67 ±0.04	0.68 ±0.03	t = -0.29, p = 0.78
L Fornix stria terminalis	0.42 ±0.04	0.42 ±0.03	t = 0.19, p = 0.85	0.72 ±0.06	0.70 ±0.05	t = 0.83, p = 0.41
R Sup long fasciculus	0.38 ±0.04	0.37 ±0.04	t = 0.56, p = 0.58	0.81 ±0.08	0.83 ±0.09	t = -0.91, p = 0.37
L Sup long fasciculus	0.34 ±0.04	0.36 ±0.02	t = -1.37, p = 0.18	1.48 ±0.14	1.43 ±0.13	t = 1.18, p = 0.25
R Sup fronto occ fasciculus	0.27 ±0.03	0.27 ±0.02	t = -0.69, p = 0.49	1.75 ±0.14	1.74 ±0.12	t = 0.29, p = 0.77
L Sup fronto occ fasciculus	0.63 ±0.04	0.64 ±0.02	t = -0.81, p = 0.43	0.77 ±0.03	0.77 ±0.02	t = 0.87, p = 0.39
R Uncinate fasciculus	0.32 ±0.05	0.33 ±0.04	t = -0.22, p = 0.82	1.38 ±0.15	1.31 ±0.13	t = 1.52, p = 0.14
L Uncinate fasciculus	0.45 ±0.04	0.45 ±0.02	t = -0.29, p = 0.77	0.68 ±0.03	0.68 ±0.03	t = 0.67, p = 0.49
R Tapetum	0.46 ±0.04	0.46 ±0.02	t = -0.15, p = 0.88	0.66 ±0.03	0.65 ±0.02	t = 0.59, p = 0.56
L Tapetum	0.47 ±0.04	0.48 ±0.03	t = -0.85, p = 0.39	0.70 ±0.02	0.68 ±0.03	t = 1.76, p = 0.09**

\*MD value has been multiplied by 1000

\*p<0.1, \*\*p<0.05

MA + = Methamphetamine user; MA - = healthy comparison subject

FA = fractional anisotropy; MD = mean diffusion; CC = corpus callosum; WM = white matter; Sup = superior; Inf = inferior; Ant = anterior; Post = posterior; Int = internal; Ext = external; long = longitudinal; Occ = occipital

Table S2: Adjusted group differences in DTI metrics for fractional anisotropy (FA)

ROI	Group est	Group std. err	Group t value	Group p	Age est	Age std. err	Age t value	Age p	Gender est	Gender std. err	Gender t value	Gender p	Model F statistic
Middle cerebellar peduncle	0,011	0,009	t = 1,24	p = 0,22	-0,001	0,001	t = -1,26	p = 0,22	0,018	0,010	t = 1,87	<b>p = 0,07*</b>	1,53
Pontine crossing tract	0,012	0,010	t = 1,17	p = 0,25	-0,001	0,001	t = -1,14	p = 0,26	0,022	0,011	t = 2,09	<b>p = 0,04**</b>	1,69
Genu CC	0,014	0,012	t = 1,21	p = 0,23	0,000	0,001	t = -0,38	p = 0,70	0,007	0,013	t = 0,59	p = 0,56	0,53
Body CC	0,006	0,011	t = 0,48	p = 0,63	0,000	0,001	t = -0,06	p = 0,95	0,011	0,012	t = 0,92	p = 0,36	0,32
Splenium CC	0,002	0,009	t = 0,24	p = 0,81	0,001	0,001	t = 0,83	p = 0,41	0,011	0,010	t = 1,14	p = 0,26	0,94
Fornix	0,004	0,009	t = 0,50	p = 0,62	0,001	0,001	t = 1,10	p = 0,28	0,010	0,009	t = 1,03	p = 0,31	1,11
R Corticospinal tract	0,007	0,011	t = 0,66	p = 0,51	-0,001	0,001	t = -1,11	p = 0,27	0,011	0,012	t = 0,90	p = 0,37	0,58
L Corticospinal tract	0,010	0,010	t = 0,95	p = 0,35	-0,001	0,001	t = -1,29	p = 0,20	0,018	0,011	t = 1,60	p = 0,12	1,19
R Medial lemniscus	0,006	0,011	t = 0,51	p = 0,61	0,000	0,001	t = -0,41	p = 0,68	0,018	0,012	t = 1,53	p = 0,14	0,78
L Medial lemniscus	0,009	0,009	t = 1,04	p = 0,30	-0,001	0,001	t = -0,61	p = 0,55	0,018	0,010	t = 1,87	<b>p = 0,07*</b>	1,29
R Inf cerebellar peduncle	0,002	0,008	t = 0,27	p = 0,79	-0,001	0,001	t = -0,83	p = 0,41	0,009	0,009	t = 1,02	p = 0,32	0,45
L Inf cerebellar peduncle	0,012	0,011	t = 1,05	p = 0,30	-0,001	0,001	t = -0,65	p = 0,52	0,003	0,012	t = 0,24	p = 0,81	0,46
R Sup cerebellar peduncle	0,005	0,007	t = 0,71	p = 0,48	-0,001	0,001	t = -0,80	p = 0,43	0,013	0,008	t = 1,70	p = 0,10	1,02
L Sup cerebellar peduncle	-0,002	0,012	t = -0,14	p = 0,89	-0,001	0,001	t = -1,29	p = 0,21	-0,001	0,013	t = -0,06	p = 0,95	0,64
R Cerebral peduncle	0,010	0,012	t = 0,84	p = 0,41	-0,001	0,001	t = -0,48	p = 0,63	0,010	0,012	t = 0,81	p = 0,42	0,38
L Cerebral peduncle	0,005	0,010	t = 0,49	p = 0,63	-0,001	0,001	t = -1,25	p = 0,22	0,022	0,010	t = 2,10	<b>p = 0,04**</b>	1,62
R Ant limb of int capsule	0,003	0,009	t = 0,36	p = 0,72	-0,002	0,001	t = -2,11	<b>p = 0,04**</b>	0,017	0,009	t = 1,83	<b>p = 0,08*</b>	2,03
L Ant limb of int capsule	0,009	0,008	t = 1,11	p = 0,27	-0,001	0,001	t = -1,04	p = 0,30	0,007	0,009	t = 0,87	p = 0,39	0,73
R Post limb of int capsule	0,005	0,008	t = 0,68	p = 0,50	-0,001	0,001	t = -1,53	p = 0,13	0,012	0,008	t = 1,47	p = 0,15	1,18
L Post limb of int capsule	0,011	0,010	t = 1,18	p = 0,25	-0,002	0,001	t = -1,76	<b>p = 0,09*</b>	0,013	0,010	t = 1,26	p = 0,22	1,44
R Retrolenticular int capsule	0,010	0,009	t = 1,09	p = 0,28	-0,002	0,001	t = -2,14	<b>p = 0,04**</b>	0,009	0,009	t = 0,99	p = 0,33	1,75
L Retrolenticular int capsule	-0,002	0,011	t = -0,22	p = 0,82	-0,001	0,001	t = -0,74	p = 0,47	-0,002	0,012	t = -0,20	p = 0,84	0,26
R Ant corona radiata	0,009	0,012	t = 0,78	p = 0,44	-0,001	0,001	t = -1,05	p = 0,30	0,013	0,012	t = 1,09	p = 0,28	0,66
L Ant corona radiata	0,005	0,010	t = 0,53	p = 0,60	-0,001	0,001	t = -1,35	p = 0,19	0,000	0,011	t = 0,05	p = 0,96	0,71
R Sup corona radiata	-0,003	0,011	t = -0,31	p = 0,76	-0,001	0,001	t = -0,89	p = 0,38	0,003	0,011	t = 0,26	p = 0,79	0,33
L Sup corona radiata	0,002	0,009	t = 0,21	p = 0,84	-0,001	0,001	t = -0,82	p = 0,42	0,011	0,009	t = 1,17	p = 0,25	0,54
R Post corona radiata	0,004	0,007	t = 0,56	p = 0,58	-0,001	0,001	t = -1,23	p = 0,23	0,016	0,008	t = 2,13	<b>p = 0,04**</b>	1,64
L Post corona radiata	0,004	0,008	t = 0,59	p = 0,56	-0,001	0,001	t = -0,76	p = 0,46	0,011	0,008	t = 1,41	p = 0,17	0,72
R Post thalamic radiation	-0,002	0,011	t = -0,17	p = 0,87	0,000	0,001	t = 0,12	p = 0,90	0,020	0,012	t = 1,72	<b>p = 0,09*</b>	1,24
L Post thalamic radiation	0,002	0,011	t = 0,19	p = 0,85	0,000	0,001	t = -0,07	p = 0,94	0,021	0,012	t = 1,74	<b>p = 0,09*</b>	1,09
R Sagittal stratum	-0,002	0,016	t = -0,16	p = 0,88	0,000	0,002	t = 0,27	p = 0,79	-0,012	0,017	t = -0,69	p = 0,49	0,16
L Sagittal stratum	0,008	0,015	t = 0,54	p = 0,59	-0,001	0,001	t = -0,65	p = 0,52	0,014	0,016	t = 0,91	p = 0,37	0,36
R Ext capsule	0,005	0,013	t = 0,38	p = 0,70	-0,001	0,001	t = -0,72	p = 0,48	-0,005	0,014	t = -0,35	p = 0,73	0,35
L Ext capsule	0,028	0,017	t = 1,69	p = 0,10	-0,002	0,002	t = -1,25	p = 0,22	0,020	0,018	t = 1,10	p = 0,28	1,37
R Cingulum	0,020	0,012	t = 1,61	p = 0,12	-0,001	0,001	t = -0,48	p = 0,63	0,011	0,013	t = 0,80	p = 0,43	0,93
L Cingulum	0,004	0,009	t = 0,49	p = 0,63	-0,001	0,001	t = -0,83	p = 0,41	0,010	0,009	t = 1,02	p = 0,31	0,47
R Cingulum hippocampus	0,008	0,007	t = 1,09	p = 0,28	0,000	0,001	t = -0,68	p = 0,50	0,008	0,007	t = 1,01	p = 0,32	0,63
L Cingulum hippocampus	0,014	0,013	t = 1,10	p = 0,28	-0,001	0,001	t = -0,82	p = 0,42	0,022	0,014	t = 1,56	p = 0,13	1,03
R Fornix stria terminalis	0,006	0,013	t = 0,45	p = 0,65	-0,001	0,001	t = -0,57	p = 0,57	0,016	0,014	t = 1,17	p = 0,25	0,48
L Fornix stria terminalis	0,000	0,011	t = 0,02	p = 0,98	0,001	0,001	t = 0,52	p = 0,60	0,012	0,012	t = 1,05	p = 0,30	0,63
R Sup long fasciculus	-0,009	0,013	t = -0,73	p = 0,47	0,000	0,001	t = -0,24	p = 0,81	-0,013	0,014	t = -0,94	p = 0,36	0,49
L Sup long fasciculus	0,019	0,010	t = 1,96	<b>p = 0,06*</b>	-0,002	0,001	t = -1,94	<b>p = 0,06*</b>	0,021	0,010	t = 2,07	<b>p = 0,05*</b>	2,72
R Sup fronto occ fasciculus	0,007	0,007	t = 0,97	p = 0,34	0,000	0,001	t = -0,41	p = 0,68	0,010	0,008	t = 1,25	p = 0,22	0,68
L Sup fronto occ fasciculus	0,011	0,011	t = 1,08	p = 0,29	-0,001	0,001	t = -1,10	p = 0,28	0,012	0,011	t = 1,09	p = 0,29	0,82

R Uncinate fasciculus	0,006	0,015	t = 0,37	p = 0,71	-0,003	0,001	t = -2,10	<b>p = 0,04**</b>	0,001	0,016	t = 0,03	p = 0,97	1,61
L Uncinate fasciculus	0,007	0,010	t = 0,66	p = 0,51	-0,001	0,001	t = -1,01	p = 0,32	0,015	0,011	t = 1,38	p = 0,18	0,79
R Tapetum	0,006	0,011	t = 0,54	p = 0,59	-0,001	0,001	t = -0,54	p = 0,59	0,018	0,011	t = 1,55	p = 0,13	0,82
L Tapetum	0,013	0,011	t = 1,26	p = 0,22	-0,001	0,001	t = -0,68	p = 0,50	0,020	0,011	t = 1,70	p = 0,10	1,22

\*p<0.1, \*\*p<0.05

MA + = Methamphetamine user; MA - = healthy comparison subject; ROI = region of interest

FA = fractional anisotropy; MD = mean diffusion; CC = corpus callosum; WM = white matter; Sup = superior; Inf = inferior; Ant = anterior; Post = posterior; Int = internal; Ext = external; long = longitudinal; Occ = occipital

Table S3: Adjusted group differences in DTI metrics for mean diffusion (MD)

ROI	Group est * 10000	Group std err * 10000	Group t value	Group p	Age est * 10000	Age std err * 10000	Age t value	Age p	Gender est * 10000	Gender std err * 10000	Gender t value	Gender p	Model F-statistic
Middle cerebellar peduncle	0,0206	0,0994	t = 0,21	p = 0,84	-0,0107	0,0096	t = -1,12	p = 0,27	0,0232	0,1060	t = 0,22	p = 0,83	0,42
Pontine crossing tract	-0,0642	0,0625	t = -1,03	p = 0,31	-0,0064	0,0060	t = -1,06	p = 0,29	-0,0202	0,0668	t = -0,30	p = 0,76	0,89
Genu CC	0,0507	0,1030	t = 0,49	p = 0,62	-0,0224	0,0099	t = -2,25	p = <b>0,03**</b>	0,0535	0,1100	t = 0,49	p = 0,63	1,72
Body CC	0,0556	0,0955	t = 0,58	p = 0,56	-0,0214	0,0092	t = -2,31	p = <b>0,03**</b>	0,0048	0,1020	t = 0,05	p = 0,96	1,99
Splenium CC	0,0246	0,1730	t = 0,14	p = 0,89	-0,0258	0,0167	t = -1,55	p = 0,13	0,0857	0,1850	t = 0,46	p = 0,65	0,79
Fornix	0,0517	0,2070	t = 0,25	p = 0,80	-0,0342	0,0200	t = -1,71	p = <b>0,09*</b>	0,2090	0,2220	t = 0,94	p = 0,35	1,04
R Corticospinal tract	-0,0847	0,0849	t = -0,99	p = 0,33	-0,0050	0,0082	t = -0,61	p = 0,55	-0,0062	0,0908	t = -0,07	p = 0,95	0,52
L Corticospinal tract	-0,0636	0,0734	t = -0,87	p = 0,39	-0,0044	0,0071	t = -0,61	p = 0,54	-0,1030	0,0785	t = -1,31	p = 0,19	1,07
R Medial lemniscus	-0,0829	0,0995	t = -0,83	p = 0,41	0,0022	0,0096	t = 0,23	p = 0,82	-0,3110	0,1060	t = -2,92	p = <b>0,01**</b>	2,99
L Medial lemniscus	0,0477	0,0847	t = 0,56	p = 0,58	-0,0069	0,0082	t = -0,84	p = 0,40	-0,0810	0,0906	t = -0,89	p = 0,38	0,89
R Inf cerebellar peduncle	-0,0408	0,0861	t = -0,47	p = 0,64	0,0085	0,0083	t = 1,01	p = 0,32	-0,0636	0,0921	t = -0,69	p = 0,49	0,42
L Inf cerebellar peduncle	-0,0268	0,0894	t = -0,30	p = 0,77	-0,0095	0,0086	t = -1,09	p = 0,28	-0,0127	0,0956	t = -0,13	p = 0,89	0,52
R Sup cerebellar peduncle	0,0221	0,0445	t = 0,49	p = 0,62	-0,0010	0,0043	t = -0,23	p = 0,82	-0,1790	0,0476	t = -3,77	p = <b>0,001**</b>	6,03
L Sup cerebellar peduncle	-0,0621	0,0888	t = -0,69	p = 0,49	0,0192	0,0086	t = 2,24	p = <b>0,03**</b>	-0,0710	0,0949	t = -0,75	p = 0,46	1,72
R Cerebral peduncle	-0,0424	0,0967	t = -0,44	p = 0,66	0,0039	0,0094	t = 0,42	p = 0,68	-0,1600	0,1030	t = -1,54	p = 0,13	0,79
L Cerebral peduncle	-0,0013	0,0928	t = -0,01	p = 0,99	0,0117	0,0089	t = 1,30	p = 0,20	-0,3060	0,0992	t = -3,08	p = <b>0,003**</b>	3,44
R Ant limb of int capsule	0,0349	0,0919	t = 0,38	p = 0,71	0,0085	0,0089	t = 0,96	p = 0,34	-0,1050	0,0983	t = -1,06	p = 0,29	0,70
L Ant limb of int capsule	-0,0709	0,0873	t = -0,81	p = 0,42	0,0243	0,0084	t = 2,88	p = <b>0,01**</b>	-0,1340	0,0934	t = -1,43	p = 0,16	2,92
R Post limb of int capsule	0,0285	0,0658	t = 0,43	p = 0,67	0,0193	0,0064	t = 3,03	p = <b>0,004**</b>	-0,1820	0,0703	t = -2,59	p = <b>0,01**</b>	4,66
L Post limb of int capsule	-0,0662	0,1110	t = -0,59	p = 0,56	0,0338	0,0108	t = 3,14	p = <b>0,003**</b>	-0,1190	0,1190	t = -0,99	p = 0,33	3,29
R Retrolenticular int capsule	-0,0556	0,1140	t = -0,49	p = 0,63	0,0253	0,0110	t = 2,30	p = <b>0,02**</b>	-0,0062	0,1220	t = -0,05	p = 0,96	1,95
L Retrolenticular int capsule	-0,1610	0,1500	t = -1,08	p = 0,29	0,0285	0,0145	t = 1,97	p = <b>0,06*</b>	0,1040	0,1610	t = 0,65	p = 0,52	2,26
R Ant corona radiata	0,0306	0,2380	t = 0,13	p = 0,89	0,0147	0,0230	t = 0,64	p = 0,53	0,0627	0,2540	t = 0,25	p = 0,81	0,21
L Ant corona radiata	-0,0584	0,1570	t = -0,37	p = 0,71	0,0313	0,0152	t = 2,06	p = <b>0,05*</b>	0,0300	0,1680	t = 0,18	p = 0,86	1,67
R Sup corona radiata	-0,1630	0,1120	t = -1,46	p = 0,15	0,0198	0,0108	t = 1,83	p = <b>0,08*</b>	-0,1490	0,1200	t = -1,25	p = 0,22	1,69
L Sup corona radiata	-0,1170	0,1180	t = -0,99	p = 0,32	0,0148	0,0114	t = 1,30	p = 0,20	-0,1260	0,1260	t = -0,10	p = 0,92	0,87
R Post corona radiata	-0,0610	0,0853	t = -0,71	p = 0,48	0,0131	0,0083	t = 1,59	p = 0,12	-0,2040	0,0912	t = -2,23	p = <b>0,03**</b>	1,98
L Post corona radiata	0,0885	0,0835	t = 1,06	p = 0,29	0,0064	0,0081	t = 0,79	p = 0,43	-0,0988	0,0893	t = -1,11	p = 0,28	1,15
R Post thalamic radiation	-0,0933	0,0880	t = -1,06	p = 0,29	0,0080	0,0085	t = 0,94	p = 0,36	-0,1260	0,0941	t = -1,34	p = 0,19	0,88
L Post thalamic radiation	-0,0061	0,0771	t = -0,08	p = 0,94	0,0080	0,0075	t = 1,07	p = 0,29	-0,0586	0,0824	t = -0,71	p = 0,48	0,45
R Sagittal stratum	0,0927	0,1430	t = 0,65	p = 0,52	-0,0093	0,0138	t = -0,67	p = 0,51	0,2720	0,1530	t = 1,78	p = <b>0,08*</b>	1,07
L Sagittal stratum	-0,0733	0,1760	t = -0,42	p = 0,68	-0,0010	0,0171	t = -0,05	p = 0,96	0,0151	0,1890	t = 0,08	p = 0,94	0,07
R Ext capsule	-0,2620	0,1120	t = -2,33	p = <b>0,03**</b>	0,0116	0,0108	t = 1,07	p = 0,29	-0,0192	0,1200	t = -0,16	p = 0,87	2,15
L Ext capsule	-0,2870	0,1760	t = -1,63	p = 0,11	0,0401	0,0170	t = 2,35	p = <b>0,02**</b>	-0,1560	0,1890	t = -0,83	p = 0,41	2,43
R Cingulum	-0,0478	0,0906	t = -0,53	p = 0,60	0,0067	0,0088	t = 0,76	p = 0,45	-0,0815	0,0969	t = -0,84	p = 0,41	0,36
L Cingulum	-0,0571	0,0817	t = -0,69	p = 0,49	0,0200	0,0079	t = 2,53	p = <b>0,02**</b>	-0,1020	0,0873	t = -1,16	p = 0,25	2,22
R Cingulum hippocampus	0,0394	0,0663	t = 0,59	p = 0,56	0,0111	0,0064	t = 1,74	p = <b>0,09*</b>	-0,0454	0,0709	t = -0,64	p = 0,53	1,27
L Cingulum hippocampus	-0,0860	0,0984	t = -0,87	p = 0,39	0,0079	0,0095	t = 0,84	p = 0,41	-0,1100	0,1050	t = -1,04	p = 0,30	0,58
R Fornix stria terminalis	-0,0160	0,1200	t = -0,13	p = 0,89	0,0133	0,0116	t = 1,15	p = 0,26	-0,1850	0,1280	t = -1,45	p = 0,16	0,912
L Fornix stria terminalis	-0,2050	0,1710	t = -1,20	p = 0,24	-0,0083	0,0165	t = -0,50	p = 0,62	-0,3370	0,1830	t = -1,84	p = <b>0,07*</b>	1,77
R Sup long fasciculus	0,3680	0,2680	t = 1,37	p = 0,18	-0,0169	0,0259	t = -0,65	p = 0,52	0,5410	0,2870	t = 1,89	p = <b>0,07*</b>	1,47
L Sup long fasciculus	-0,6300	0,4350	t = -1,45	p = 0,16	0,0723	0,0420	t = 1,72	p = 0,09	-0,3690	0,4650	t = -0,79	p = 0,43	1,49
R Sup fronto occ fasciculus	-0,2220	0,4400	t = -0,50	p = 0,63	0,0189	0,0425	t = 0,44	p = 0,66	-0,4010	0,4700	t = -0,85	p = 0,39	0,28

L Sup fronto occ fasciculus	-0,0984	0,0959	t = -1,03	p = 0,31	0,0182	0,0093	t = 1,96	p = <b>0,06*</b>	-0,0183	0,1030	t = -0,18	0,86	1,60
R Uncinate fasciculus	-0,7590	0,4250	t = -1,79	p = <b>0,08*</b>	0,1080	0,0410	t = 2,64	p = <b>0,01*</b>	-0,1060	0,4540	t = -0,23	0,82	3,29
L Uncinate fasciculus	-0,0270	0,1050	t = -0,26	p = 0,79	-0,0105	0,0101	t = -1,04	p = 0,31	0,1640	0,1120	t = 1,47	0,15	1,01
R Tapetum	-0,0222	0,0887	t = -0,25	p = 0,80	-0,0034	0,0086	t = -0,41	p = 0,69	0,1200	0,0948	t = 1,26	0,21	0,65
L Tapetum	-0,1640	0,0857	t = -1,91	p = <b>0,06*</b>	0,0001	0,0083	t = 0,01	p = 0,99	-0,0886	0,0917	t = -0,97	0,34	1,36

\*p<0.1, \*\*p<0.05

MA + = Methamphetamine user; MA - = healthy comparison subject; ROI = region of interest

## **APPENDICES**

- A. HREC approval letter 263/2010**
- B. HREC approval letter 635/2014**
- C. MATRIX information session (slides): 28 June 2011**
- D. Candidate approval for MPhil (Neuropsychiatry) UCT**
- E. City of Cape Town permission to recruit from Matrix Clinic**
- F. Inclusion/exclusion criteria flyer for Matrix and Mitchells Plain  
Community Health Centre**
- G. Participant informed consent sheet**
- H. Author Guidelines**

## APPENDIX A



UNIVERSITY OF CAPE TOWN

Health Sciences Faculty  
Research Ethics Committee  
Room E52-24 Groote Schuur Hospital Old Main Building  
Observatory 7925  
Telephone [021] 406 6626 • Facsimile [021] 406 6411  
e-mail: shuretta.thomas@uct.ac.za

06 July 2010

HREC REF: 263/2010

Dr H Gouse  
GSH-HIV mental Health Group  
J2, Office 76  
GSH

Dear Dr Gouse

**PROJECT TITLE: METHAMPHETAMINE INDUCED DEFICITS IN NEURO-AND-SOCIAL-  
COGNITION IN THE CONTEXT OF HIV INFECTION.**

Thank you for addressing the issues raised by the Faculty of Health Sciences Human Research Ethics Committee.

It is a pleasure to inform you that the Ethics Committee has **formally approved** the above-mentioned study.

**Approval is granted for one year till the 15<sup>th</sup> July 2011.**

Please submit an annual progress report if the research continues beyond the expiry date. Please submit a brief summary of findings if you complete the study within the approval period so that we can close our file.

Please note that the ongoing ethical conduct of the study remains the responsibility of the principal investigator.

**Please quote the REC. REF in all your correspondence.**

Yours sincerely



Signature removed

 **PROFESSOR M BLOCKMAN**  
**CHAIRPERSON, HSF HUMAN ETHICS**  
Federal Wide Assurance Number: FWA00001637.

S Thomas



## APPENDIX B



**UNIVERSITY OF CAPE TOWN**  
**Faculty of Health Sciences**  
**Human Research Ethics Committee**

Room E52-24 Old Main Building  
Groote Schuur Hospital  
Observatory 7925  
Telephone [021] 406 6338 • Facsimile [021] 406 6411  
Email: shuretta.thomas@uct.ac.za  
Website: [www.health.uct.ac.za/fhs/research/humanethics/forms](http://www.health.uct.ac.za/fhs/research/humanethics/forms)

---

29 August 2014

**HREC REF: 635/2014**

**A/Prof John Joska**  
Psychiatry & Mental Health  
J-Block, GSH

Dear A/Prof Joska

**PROJECT TITLE: WHITE MATTER CORRELATES OF NEUROPSYCHOLOGICAL FUNCTION IN YOUNG, METHAMPHETAMINE DEPENDENT INDIVIDUALS - A DIFFUSION TENSOR IMAGING STUDY - SUB-STUDY LINKED TO 236/2010 (MPHIL CANDIDATE - DR C FREEMAN)**

Thank you for submitting your sub-study to the Faculty of Health Sciences Human Research Ethics Committee for review.

It is a pleasure to inform you that the HREC has **formally approved** the above-mentioned study.

**Approval is granted for one year until the 30<sup>th</sup> August 2015.**

Please submit a progress form, using the standardised Annual Report Form if the study continues beyond the approval period. Please submit a Standard Closure form if the study is completed within the approval period.  
(Forms can be found on our website: [www.health.uct.ac.za/fhs/research/humanethics/forms](http://www.health.uct.ac.za/fhs/research/humanethics/forms))

***We acknowledge that the MPHIL student, Dr Carla Patricia Freeman will also be involved in this study.***

Please note that the ongoing ethical conduct of the study remains the responsibility of the principal investigator.

**Please quote the HREC reference no in all your correspondence.**

Yours sincerely

signature removed

PP **PROFESSOR M BLOCKMAN**  
**CHAIRPERSON, FHS HUMAN ETHICS**  
Federal Wide Assurance Number: FWA00001637.  
Institutional Review Board (IRB) number: IRB00001938

HREC 635/2014

## APPENDIX C

19/05/2016

### What does my brain look like?

Matrix clinic, Cape Town.  
28<sup>th</sup> June 2011  
Dr. Carla Freeman

### Magnetic Resonance Image MRI



### How does MRI work?

- "The MAGNET"
- A powerful magnetic field is created in the bore
- Atoms are lined up
- Photographic "slices" of the brain can be seen
- Taken a step further, these images can be taken while performing a task

### MRI and Safety




- No radiation
- Body returns to normal
- No long term damage
- Caution with metal objects



### The sexiest organ in the body!





### The MRI equivalent




Not all brains are equal...




A quiz...




An example of an abnormal scan



How does tik affect the brain



### Methamphetamine and the brain...



### How does tik affect the body?



### "Meth mouth"



### Time to reversal of the damage?

- Up to one year to reverse some effects
- The longer in a structured rehab, the better the outcome
- Longer use had worse outcomes
- RECOVERY IS DIFFICULT BUT IT IS POSSIBLE

Questions?

## APPENDIX D

### Re: Freeman: Confirmation of Approval of Study Proposal

John Joska <john.joska@uct.ac.za>

Mon 2015-01-26 05:04 PM

Inbox

To: Carla Freeman <FRMCAR002@myuct.ac.za>;

The whip is ever on your back...!

---

**From:** Vuyi Mgoqi <vuyi.mgoqi@uct.ac.za>

**Date:** Monday 26 January 2015 at 5:02 PM

**To:** Carla Freeman <FRMCAR002@myuct.ac.za>

**Cc:** Jackie Cogill <jackie.cogill@uct.ac.za>, John Joska <john.joska@uct.ac.za>

**Subject:** Freeman: Confirmation of Approval of Study Proposal

Dear Dr Freeman


#### Candidature Approval (FRMCAR002)

Degree	MPhil in Neuropsychiatry
Title	White matter correlates of neuropsychological function in young methamphetamine dependent individuals – a diffusion tensor imaging study
Department	Psychiatry and Mental Health
Supervisor	A/Prof J Joska
Ethics Approval	635/2014

I am pleased to advise that the Chair of the Dissertations/Doctoral & Masters Committee has approved your candidature for the above degree on behalf of the Committee. Formal approval was obtained by publication in the Dean's Circular, PG-Med Nov-Dec2014.

Yours sincerely

Vuyi Mgoqi

 Vuyiseka Mgoqi   Receptionist: PG Academic Administration   Faculty of Health Sciences   University of Cape Town   Room N2.19, Wernher & Beit North, Health Sciences Campus, Anzio Rd, Observatory, 7925   (+ 2721 406 67516 + 27 21 406 6584   Office Hours: 08h30 - 16h30 Unavailable Hours: 13h00 - 13h30
--

## APPENDIX E



Civic Centre  
12 Hertzog Boulevard  
Cape Town 8001  
P O Box 2815, Cape Town 8000  
**Ask for: Dr G H Visser**  
Tel: 021 400-3981  
Cell: 083 298 8718  
Fax: 021 421-4894  
E-mail: [helene.visser@capetown.gov.za](mailto:helene.visser@capetown.gov.za)  
Website: <http://www.capetown.gov.za>  
Ref:  
Filename: G:\Research\Gouws 10207.docx

Iziko iLuntu  
12 Hertzog Boulevard  
Cape Town 8001  
P O Box 2815, Cape Town 8000  
**Cela: Qrh G H Visser**  
Umxweba: 021 400-3981  
Cell: 083 298 8718  
Iheksi: 021 421-4894

Burgersentrum  
Hertzog-boulevard 12  
Kaapstad 8001  
Postbus 2815, Kaapstad 8000  
**Vra vir: Dr G H Visser**  
Tel: 021 400-3981  
Sel: 083 298 8718  
Faks: 021 421-4894

CITY HEALTH — Specialised Health  
2010-11-22

Dear Dr Gouws

**re: Research Request: Methamphetamine induced Deficits in Neuro-and-Social cognition in the Context of HIV Infection (ID: 10207)**

Permission has been granted to do the research as per your protocol at the following City Health Matrix Sites (which include the required recruitment of HIV positive & control clients from the same clinics), Tafelsig, Table View and Delft South.

Your request to also recruit clients from adjoining ARV sites was not approved due to the following complexities: Tafelsig and Table View don't have ARV clinics and the nearest ARV site would be Mzamomhle and Du Noon respectively. However, these 2 sites are very busy, space constrained sites, not suitable for research recruitment.

The Health Management Team therefore recommended that you start with the Matrix Sites and approach us again at a later stage if enrolment is too slow and we could consider other sites to recruit the HIV positive clients from.

**Please note the following:**

1. Any individual patient information obtained must be kept confidential.
2. Access to the facilities must be arranged with the relevant Managers such that normal activities are not disrupted.
3. A copy of your final report should be sent to the City Health Head Office, P O Box 2815 Cape Town 8001, within 3 months of its completion and feedback must also be given to the clinics involved.
4. Your project has been given an ID Number (10207). Please use this in any future correspondence with us.

Thank you for your cooperation and please contact me if you require further information or assistance.

Yours sincerely

signature removed

DR G H VISSER  
**MANAGER: SPECIALISED HEALTH**

cc. Mrs Elloker & Mrs Nqana  
Mrs Sifanelo & Mrs Stanley  
Mr Nkoko & Mrs Nojaholo  
Dr Jennings  
Mrs Bosch

THIS CITY WORKS FOR YOU ESI SIXEKO SISEBENZELA WENA HIERDIE STAD WERK VIR JOU



## **APPENDIX F**

1<sup>st</sup> Floor  
Anzio Road  
Groote Schuur Hospital  
Observatory  
Cape Town  
7925

Tel: +27 21 404 5222  
Fax: +27 21 448 8158  
Email:  
Dr Hetta Gouse  
[hetta.gouse@uct.ac.za](mailto:hetta.gouse@uct.ac.za)  
Dr Carla Freeman  
[carla.freeman@uct.ac.za](mailto:carla.freeman@uct.ac.za)  
Website: [www.hivmentalhealth.co.za](http://www.hivmentalhealth.co.za)

**THIS PATIENT IS PART OF THE GSH-  
HIV MENTAL HEALTH GROUP  
RESEARCH PROJECT.  
AN MRI SCAN IS AVAILABLE ON  
REQUEST.**

### **Inclusion criteria**

- a) Between 18 and 40 years old.
- b) Conversational in English.
- c) HIV+ HAART naïve with CD4 cell count of  $\leq 350$ .
- d) Literate with minimum grade 7 (Std 5) education.

### **Exclusion criteria**

- a) Past and present comorbid psychiatric disorders.
- b) Past and present comorbid neurological disorders.
- c) Traumatic head injury with loss of consciousness for longer than 30 minutes, AND/OR requiring overnight admission to hospital.
- d) Contra-indications to MRI – such as pregnancy, metal or claustrophobia.

## **APPENDIX G**

### **MATRIX PARTICIPANT INFORMATION LEAFLET AND CONSENT FORM**

#### **Methamphetamine induced deficits in neuro- and social-cognition in the context of HIV infection.**

PRINCIPAL INVESTIGATOR: Dr John Joska      CONTACT NUMBER: 021-404 2154  
ADDRESS: Department of Psychiatry and Mental Health, J-block, Groote Schuur Hospital,  
Anzio Road, Observatory, 7925

As you enter Matrix treatment some information will be collected from you by the clinic staff. You will also be required to do an HIV test. We would like to use some of this information, including your HIV status, to help us understand drug addiction better and request your permission for access to this information. After this you may be contacted to be screened to see if you meet the criteria to partake in the rest of the research study which consists of two parts, neuropsychological testing and a MRI scan. You will be interviewed in order to find out if you have any mental problems which may affect your future health or your ability to function.

During the first visit at Groote Schuur Hospital you will perform a series of tests in order to see how well you can think, remember and do certain things. It will take about 3 hours. This will be followed by a medical examination performed by a medical doctor. Voluntary pre and post-test counselling will accompany a blood test to confirm your HIV status. These results will remain strictly confidential. Two additional tubes of blood will be taken for genetic testing. This blood will NOT be used to diagnose genetic illnesses. It will only be used to further understand how drugs like methamphetamine and illnesses like HIV affect the brain. These results will be kept confidential and will not impact on your current or future care and treatment. This procedure involves one needle-prick and we will do our utmost to ensure your comfort throughout.

We would like to use this information, as well as basic information about your medical condition to study the problems that people with HIV/AIDS AND/OR tik use (methamphetamine) might have. It will be kept private and not linked to your name. If your doctor or Matrix Clinic staff requests it, a report on your condition will be given to your clinic doctor or counsellor who referred you. They will use this to make a decision on your treatment; furthermore it may lead to better treatment for others in the future.

This study may make you feel uncomfortable as you talk about mental health problems. You may feel embarrassed or shy. Sometimes painful information is shared. Also, some people feel that it is better not to know about memory or thinking problems.

During the second visit in this study you will have a type of brain scan which will be done at the Cape Universities Brain Imaging Centre at Tygerberg Hospital. This scan is called an MRI (magnetic resonance imaging) scan. The scan will require you to lie on your back on a table that will move into the scanning machine for the 60 minutes it will take for the scan. During most of this time you will be able to close your eyes and rest. You will also be able to talk to the study doctor/assistant at all times during the scan if you should experience any discomfort. You will do two tasks during which you will be asked to push a button. The scan is a safe procedure if you have been screened



correctly for the presence of any magnetic material on or inside you such as pace-makers, surgical clips and metal objects in the eyes. A formal screen for this will be done at the screening visit by a member of the study team. As the scan is done in a relatively confined space, occasionally people become anxious. This does not happen often, and if you feel anxious, we will spend time allowing you to get used to the surroundings before we begin. When the magnet in the machine is switched on, it will make some loud banging noises, but you will be clearly warned when this will take place. At this time you will feel nothing and the noise is not harmful to you in any way. To minimise the possible discomfort associated with this, we will give you some soft earplugs and will also put earphones on so that you can listen to music if you so choose.

These are the main risks. You should feel free to mention your feelings or concerns to any member of the study team.

You should feel free not participate or to not answer any question you are uncomfortable with. We would like you to sign this page to say that you allow us to use this information, but you are free NOT to. This will not affect your care and treatment in any way. It remains between you and the interviewer.

#### **Methamphetamine users only**

The Matrix clinic staff requested that we make some of the information that we collect available to them. We are going to ask you some personal questions, including questions about your sexual behaviour over the last four months. Should you not want this or any other information to be made available to the Matrix clinic staff please indicate it by ticking the box below We would like to contact you in six months for a follow up study.

☐ DO NOT MAKE MY INFORMATION AVAILABLE TO MATRIX CLINIC STAFF.

**This study has been approved by the Human Research Ethics Committee (HREC) of the Faculty of Health Sciences of the University of Cape Town** and will be conducted according to the ethical guidelines and principles of the international Declaration of Helsinki, South African Guidelines for Good Clinical Practice and the Medical Research Council (MRC) Ethical Guidelines for Research. **For any queries regarding your rights as a participant, please contact the investigator, Dr Hetta Gouse (021 404 5224), Dr John Joska (021 404-2154) or the HREC at 021-4066411.**

Please answer the questions as honestly as possible. Ask for help if you need it.

**Patient:** "I understand what this interview is about and agree that the information may be used to study mental problems in people with HIV/AIDS"

Signed at (*place*) ..... on (*date*) .....20\_\_.

.....  
**Signature of participant**

.....  
**Signature of witness**

**Investigator/nurse/counsellor:** "I have explained this study to the patient in a way that they understand and allowed them to ask questions  
I (*name*) ..... declare that:

.....  
**Signature of investigator**

.....  
**Signature of witness**

## APPENDIX H



### PSYCHIATRY RESEARCH: NEUROIMAGING

The Official Publication of the [International Society for Neuroimaging in Psychiatry](#)

#### AUTHOR INFORMATION PACK

#### TABLE OF CONTENTS

• <b>Description</b>	<b>p.1</b>
• <b>Audience</b>	<b>p.1</b>
• <b>Impact Factor</b>	<b>p.1</b>
• <b>Abstracting and Indexing</b>	<b>p.1</b>
• <b>Editorial Board</b>	<b>p.2</b>
• <b>Guide for Authors</b>	<b>p.3</b>



ISSN: 0925-4927

#### DESCRIPTION

The *Neuroimaging* section of *Psychiatry Research* publishes manuscripts on positron emission tomography, magnetic resonance imaging, computerized electroencephalographic topography, regional cerebral blood flow, computed tomography, magnetoencephalography, autoradiography, post-mortem regional analyses, and other **imaging techniques**. Reports concerning results in **psychiatric disorders**, **dementias**, and the effects of **behaviorial tasks** and **pharmacological treatments** are featured. We also invite manuscripts on the methods of obtaining images and computer processing of the images themselves. Selected case reports are also published.

#### Benefits to authors

We also provide many author benefits, such as free PDFs, a liberal copyright policy, special discounts on Elsevier publications and much more. Please click here for more information on our [author services](#).

Please see our [Guide for Authors](#) for information on article submission. If you require any further information or help, please visit our support pages: <http://support.elsevier.com>

#### AUDIENCE

Psychiatrists, Neurologists, Psychopharmacologists and Clinical Neurophysiologists.

#### IMPACT FACTOR

2013: 2.831 © Thomson Reuters Journal Citation Reports 2014

#### ABSTRACTING AND INDEXING

BIOSIS  
Chemical Abstracts  
Current Contents/Life Sciences  
MEDLINE®  
EMBASE  
PsycINFO Psychological Abstracts  
SIIC Data Bases  
Scopus

## EDITORIAL BOARD

---

### *Editors-in-Chief*

**M.S. Buchsbaum**, San Diego, California, USA  
**T. Dierks**, Bern, Switzerland  
**K. Maurer**, Frankfurt, Germany

### *Managing Editor*

**S. Buchsbaum**

### *Associate Editors*

**N.C. Andreasen**, Iowa City, Iowa, USA  
**T.J. Crow**, Oxford, UK  
**G. Sedvall**, Stockholm, Sweden  
**D.R. Weinberger**, Baltimore, Maryland, USA

### *Editorial Board Members*

**P. Brambilla**, Udine, Italy  
**J.D. Brodie**, New York, New York, USA  
**C.S. Carter**, Sacramento, California, USA  
**L. Chang**, Honolulu, Hawaii, USA  
**P. Falkai**, Munich, Germany  
**P. Flor-Henry**, Edmonton, Alberta, Canada  
**S.A. Galderisi**, Naples, Italy  
**A. Heinz**, Berlin, Germany  
**T. Iidaka**, Nagoya, Japan  
**M. Kubicki**, Boston, Massachusetts, USA  
**V. Kumari**, London, UK  
**J.S. Kwon**, Seoul, South Korea  
**J.-L. Martinot**, Orsay, France  
**A. M. McIntosh**, Edinburgh, Scotland, UK  
**A. Meyer-Lindenberg**, Mannheim, Germany  
**V. Molina**, Salamanca, Spain  
**A.B. Newberg**, Philadelphia, Pennsylvania, USA  
**E.A. Nofzinger**, Pittsburgh, Pennsylvania, USA  
**G.D. Pearlson**, Hartford, Connecticut, USA  
**O. Pogarell**, München, Germany  
**J.D. Ragland**, Sacramento, California, USA  
**S.L. Rauch**, Belmont, Massachusetts, USA  
**M. Reite**, Denver, Colorado, USA  
**P.F. Renshaw**, Salt Lake City, Utah, USA  
**B. Saletu**, Vienna, Austria  
**F. Schneider**, Aachen, Germany  
**J. Schroeder**, Heidelberg, Germany  
**M.E. Shenton**, Boston, Massachusetts, USA  
**G.W. Small**, Los Angeles, California, USA  
**M. Suzuki**, Toyama, Japan  
**R. Szeszko**, Glen Oaks, New York, USA  
**W.D. Taylor**, Durham, North Carolina, USA  
**J. Théberge**, London, Ontario, Canada  
**N. E. M. van Haren**, Utrecht, The Netherlands  
**A. Vita**, Milan, Italy  
**L.O. Wahlund**, Huddinge, Sweden  
**M. Yücel**, Melbourne, Victoria, Australia

## GUIDE FOR AUTHORS

---

The Neuroimaging section of *Psychiatry Research* publishes manuscripts on positron emission tomography, magnetic resonance imaging, magnetic resonance spectroscopy, functional magnetic resonance imaging, single photon emission computed tomography, computerized electroencephalographic topography, regional cerebral blood flow, computed tomography, magnetoencephalography, autoradiography, post-mortem regional analyses, and other imaging techniques. Reports concerning results in psychiatric disorders, dementias and the effects of behavioral tasks and pharmacological treatments are featured. We also invite manuscripts on the methods of obtaining images and computer processing of the images themselves.

### Preparation of manuscripts

**Title page.** The Title page should include the author byline, with names of authors on the same line(s). Superscript letters (a, b, c), not numerals, should be used to key institutional affiliation (if all authors are in the same department, the superscript letter should be omitted); an asterisk should be entered to designate the corresponding author. Underneath the byline, institutional affiliations should be listed (department, institution, city, state or province (if applicable) and country. Funding information should not be included on the title page but should instead be given following the Discussion section. In an asterisked Corresponding Author footnote at the bottom of the title page, telephone/fax numbers and e-mail address of the corresponding author should be provided; e-mail addresses, if desired, may also be provided for the co-authors (or co-corresponding author, if applicable).

**Abstract.** The Abstract should be 150-200 words for full-length articles and 75 words for brief reports, summarizing the aims of the study, the methods used, the results and the major conclusions. Do not include a summary at the end of the article. Note that *Psychiatry Research: Neuroimaging* does not use the structured abstract style; do not include bold-faced headings within the abstract. The Abstract should be a single paragraph. Do not include detailed statistics or p-values in the abstract; simply say "significant" or "non-significant." The Abstract should be followed by up to seven Key Words, which accord with the indexing

The abstract should be followed by up to seven key words should be listed which accord with the indexing conventions of Index Medicus. Note that the keywords should not duplicate words used in the title of the article, which will be automatically indexed.

**Text.** Although exceptions will be considered, manuscripts should not exceed 5000 words, and shorter manuscripts (e.g., 3000 words) are preferred. Each article should contain the following major headings: Introduction (preceded by arabic number 1.), Methods (preceded by number 2.), Results (preceded by number 3.), Discussion (preceded by number 4.), Acknowledgment (optional section following the discussion, which should not be preceded by a numeral), and References (should not be preceded by a numeral).

Subheadings should follow the numbering system used in the major heading; for example, the subheading "Subjects" within the Methods section should be flush left on a separate line and designated 2.1., the subheading "Procedures" should be designated 2.2., etc.

Lower level headings, if required, should also be numbered (e.g., "2.1.1. Patients." as a lower order heading under "2.1. Subjects."). Only the first letter of the first word of each heading should be capitalized.

The use of abbreviations within the text should be minimized, and each abbreviation, when introduced, must be defined and used consistently thereafter. Systeme International measurements should be used. For products or instruments (do not abbreviate) used in the research reported, provide the name, city and country of the supplier in parentheses. All tables and figures must be referred to in the text.

### Manuscript categories

**Articles.** Although exceptions will be considered, manuscripts should not exceed 5000 words, and shorter manuscripts (e.g., 3000 words) are preferred. Each article should contain the following major headings: Introduction (preceded by arabic number 1.), Methods (preceded by number 2.), Results (preceded by number 3.), Discussion (preceded by number 4.), Acknowledgment (optional section

following the discussion, which should not be preceded by a numeral), and References (should not be preceded by a numeral). Subheadings should follow the numbering system used in the major heading; for example, the subheading "Subjects" within the Methods section should be flush left on a separate line and designated 2.1., the subheading "Procedures" should be designated 2.2., etc. Lower level headings, if required, should also be numbered (e.g., "2.1.1. Patients." as a lower order heading under "2.1.Subjects."). Only the first letter of the first word of each heading should be capitalized.

**Brief reports.** Brief reports should not exceed 1500 words, including a 75-word abstract, 3 keywords, text, and references plus 1 table or 1 figure.

**Case reports.** Case reports will only be considered as Letters to the Editor (see following instructions ).

**Letters to the Editor.** Letters to the Editor should be 750-1000 words or less. The Letter should not include a title page, abstract or key words. Authors' names and affiliations should be listed at the end of the Letter, along with the corresponding author's email address. There should be no more than 5 references, and no tables or figures.

**Introduction.** The introduction should be brief and explain the purpose of the study; an extensive review of the literature should be avoided, but directly relevant articles by other investigators, as well as by the authors themselves, must be cited. If the manuscript includes subjects who have been included in previous reports, references should be provided and the number of subjects whose data have been included elsewhere should be specified.

**Methods.** The Methods should contain sufficient detail to enable others to repeat the procedures without studying the references directly.

**Results.** The Results should summarize the most important data, and statistical correlations should be included. Tabular data should not be duplicated in the text; important points and trends should be pointed out. The final sentence should emphasize the importance attached to the observations.

**Discussion.** The discussion should relate directly to the study being reported and give perspective to the adequacy of the materials and methods for the purpose of the study. Results should be interpreted to lend meaning to the observations. Any discrepancies with previously published results should be explained. The paper should conclude with a brief statement regarding the significance of the study.

**Acknowledgement.** The Acknowledgement section is an optional section and should also be used for grant-support information.

**Contributors.** The individual contributions of each author should be briefly summarized.

**Conflict of interest.** All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within three (3) years of beginning the work submitted that could inappropriately influence, or be perceived to influence, their work. Examples of potential conflicts of interest that should be disclosed include employment, consultancies, stock ownership (except for personal investment purposes equal to the lesser of one percent (1%) or USD 5000), honoraria, paid expert testimony, patent applications, registrations, and grants. If there are no conflicts of interest, authors should state that there are none.

**Abbreviations.** Define abbreviations at their first occurrence in the article. Abbreviations should be defined when they first occur in the abstract, in the text, and also in tables and figure legends. Once an abbreviation has been introduced in the main body of the text, it should be used throughout.

**Statistical reporting.** Statistical reporting should be complete, including at a minimum name of statistical test, test value, degrees of freedom where appropriate, and *p*-value. Italic font should be used for *n* (sample size) and statistical terms, e.g., *t*, *r*, *F*, *U*, *p*.

### **Submission of manuscripts**

Once a manuscript has successfully been submitted via the online submission system, authors may track the status of their manuscript using the online submission system (details will be provided by e-mail). If your manuscript is accepted by the journal, subsequent tracking facilities are available on Elsevier's Author Gateway, using the unique reference number provided by Elsevier and corresponding author name (details will be provided by e-mail).

Authors may send queries concerning the submission process or journal procedures to the Managing Editor (E-mail: [sherry.buchsbaum@gmail.com](mailto:sherry.buchsbaum@gmail.com)).

## **BEFORE YOU BEGIN**

### **Ethics in publishing**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

### **Human and animal rights**

If the work involves the use of animal or human subjects, the author should ensure that the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans <http://www.wma.net/en/30publications/10policies/b3/index.html>; EU Directive 2010/63/EU for animal experiments [http://ec.europa.eu/environment/chemicals/lab\\_animals/legislation\\_en.htm](http://ec.europa.eu/environment/chemicals/lab_animals/legislation_en.htm); Uniform Requirements for manuscripts submitted to Biomedical journals <http://www.icmje.org>. Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

### **Conflict of interest**

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. If there are no conflicts of interest then please state this: 'Conflicts of interest: none'. See also <http://www.elsevier.com/conflictsofinterest>. Further information and an example of a Conflict of Interest form can be found at: [http://help.elsevier.com/app/answers/detail/a\\_id/286/p/7923](http://help.elsevier.com/app/answers/detail/a_id/286/p/7923).

### **Submission declaration and verification**

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service CrossCheck <http://www.elsevier.com/editors/plagdetect>.

### **Changes to authorship**

This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts:

*Before the accepted manuscript is published in an online issue:* Requests to add or remove an author, or to rearrange the author names, must be sent to the Journal Manager from the corresponding author of the accepted manuscript and must include: (a) the reason the name should be added or removed, or the author names rearranged and (b) written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Requests that are not sent by the corresponding author will be forwarded by the Journal Manager to the corresponding author, who must follow the procedure as described above. Note that: (1) Journal Managers will inform the Journal Editors of any such requests and (2) publication of the accepted manuscript in an online issue is suspended until authorship has been agreed.

*After the accepted manuscript is published in an online issue:* Any requests to add, delete, or rearrange author names in an article published in an online issue will follow the same policies as noted above and result in a corrigendum.

### **Copyright**

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright, see <http://www.elsevier.com/copyright>). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <http://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: please consult <http://www.elsevier.com/permissions>.

For open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' (for more information see <http://www.elsevier.com/OAauthoragreement>). Permitted third party reuse of open access articles is determined by the author's choice of user license (see <http://www.elsevier.com/openaccesslicenses>).

### **Author rights**

As an author you (or your employer or institution) have certain rights to reuse your work. For more information on author rights for (a) subscription articles please see <http://www.elsevier.com/journal-authors/author-rights-and-responsibilities>; (b) for open access articles please see <http://www.elsevier.com/OAauthoragreement>.

### **Role of the funding source**

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

### **Funding body agreements and policies**

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some authors may also be reimbursed for associated publication fees. To learn more about existing agreements please visit <http://www.elsevier.com/fundingbodies>.

### **Funding body agreements and policies**

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some authors may also be reimbursed for associated publication fees. To learn more about existing agreements please visit <http://www.elsevier.com/fundingbodies>.

After acceptance, open access papers will be published under a noncommercial license. For authors requiring a commercial CC BY license, you can apply after your manuscript is accepted for publication.

### **Open access**

This journal offers authors a choice in publishing their research:

#### **Open access**

- Articles are freely available to both subscribers and the wider public with permitted reuse
- An open access publication fee is payable by authors or on their behalf e.g. by their research funder or institution

#### **Subscription**

- Articles are made available to subscribers as well as developing countries and patient groups through our universal access programs (<http://www.elsevier.com/access>).
- No open access publication fee payable by authors.

Your publication choice will have no effect on the peer review process or acceptance of submitted articles.

For open access articles, permitted third party (re)use is defined by the following Creative Commons user licenses:

*Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)*

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

The open access publication fee for this journal is **3000**, excluding taxes. Learn more about Elsevier's pricing policy: <http://www.elsevier.com/openaccesspricing>.

#### **Language (usage and editing services)**

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's WebShop (<http://webshop.elsevier.com/languageediting/>) or visit our customer support site (<http://support.elsevier.com>) for more information.

#### **Submission**

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

*Submit your article*

Please submit your article via <http://ees.elsevier.com/psyn>.

#### **Referees**

Please submit the names and institutional e-mail addresses of several potential referees. For more details, visit our [Support site](#). Note that the editor retains the sole right to decide whether or not the suggested reviewers are used.

#### **Editorial Policy**

Submitted manuscripts will be reviewed anonymously by at least two referees. Should a revised manuscript be required by the editors, the authors are requested to resubmit their revised manuscript to the journal within 6 months time. Studies on humans submitted to the journal must comply with the principles laid down in the Declaration of Helsinki (Br Med J 1964; 2: 177-178). The editors retain the right to reject papers on the grounds that, in their opinion, the ethical justification is questionable. Manuscripts may be edited to improve clarity and expression.

Manuscripts that are not published and that are not resubmitted in revised form will be destroyed within 1 year of the date of submission.

### **PREPARATION**

#### **Use of word processing software**

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the Guide to Publishing with Elsevier: <http://www.elsevier.com/guidepublication>). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

#### **Article structure**

The Abstract should be 150-200 words for full-length articles and 75 words for brief reports.



#### *Subdivision - numbered sections*

Divide your article into clearly defined and numbered sections. Subsections should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. (the abstract is not included in section numbering). Use this numbering also for internal cross-referencing: do not just refer to 'the text'. Any subsection may be given a brief heading. Each heading should appear on its own separate line.

#### **Essential title page information**

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that phone numbers (with country and area code) are provided in addition to the e-mail address and the complete postal address. Contact details must be kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

#### **Abstract**

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

#### **Graphical abstract**

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. See <http://www.elsevier.com/graphicalabstracts> for examples.

Authors can make use of Elsevier's Illustration and Enhancement service to ensure the best presentation of their images and in accordance with all technical requirements: [Illustration Service](#).

#### **Highlights**

Highlights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). See <http://www.elsevier.com/highlights> for examples.

#### **Keywords**

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

#### **Abbreviations**

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

In the abstract, define all abbreviations so that electronic searches for commonly used abbreviations or the full name can be successful. Avoid abbreviations unique to the current article so as to widen the circle of readers. We recognize that many abbreviations or acronyms may be more familiar to

the reader than the full name. However abbreviations and acronyms used by relatively few other published reports or abbreviations with several alternate meanings in data base searches should always be spelled out throughout the report.

### **Acknowledgements**

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

### **Footnotes**

Footnotes should be used sparingly. Number them consecutively throughout the article. Many word processors can build footnotes into the text, and this feature may be used. Otherwise, please indicate the position of footnotes in the text and list the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list.

### **Artwork**

#### *Electronic artwork*

##### *General points*

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the printed version.
- Submit each illustration as a separate file.

A detailed guide on electronic artwork is available on our website:

<http://www.elsevier.com/artworkinstructions>

**You are urged to visit this site; some excerpts from the detailed information are given here.**

##### *Formats*

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

##### **Please do not:**

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

##### *Color artwork*

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. For further information on the preparation of electronic artwork, please see <http://www.elsevier.com/artworkinstructions>.

Please note: Because of technical complications that can arise by converting color figures to 'gray scale' (for the printed version should you not opt for color in print) please submit in addition usable black and white versions of all the color illustrations.

### Illustration services

Elsevier's WebShop (<http://webshop.elsevier.com/illustrationservices>) offers Illustration Services to authors preparing to submit a manuscript but concerned about the quality of the images accompanying their article. Elsevier's expert illustrators can produce scientific, technical and medical-style images, as well as a full range of charts, tables and graphs. Image 'polishing' is also available, where our illustrators take your image(s) and improve them to a professional standard. Please visit the website to find out more.

### Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules.

### References

#### *Citation in text*

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

#### *Reference style*

**Literature citations.** References in the text to literature cited should be given by the surname of the author(s), followed by the year of publication in parentheses, e.g., Smith and Smith (2011) or (Allen et al., 2006; Smith, 2008a, 2008; Jones and Jones, 2013). For three or more authors, the surname of the first author followed by et al. should be used. For citations to articles with two authors, use "and" instead of ampersand (Jones and Smith, 2010). References should be arranged in alphabetical order by first author and should not be numbered. For single-authored articles, if more than one article by the same author is included, list each reference in chronological order. If both articles were published in the same year, alphabetize by the first major word of the article title and designate the first listed article as "a," the second as "b," etc. For multi-authored articles, list in alphabetical order by (1) last name of first author, (2) last name of second author, etc. If the names of all authors are identical, list in chronological order. If both authors' names and year of publication are the same, alphabetize by the first major word of the article title.

Provide the last names and first initials of all authors (do not use et al. in the reference list). Journal titles should not be abbreviated; provide the journal's full name. Do not italicize journal or book titles. Be sure that all references are complete: Journal articles should include authors, year of publication, article title, full journal name, volume number, and beginning and concluding page numbers. Book chapters should include authors, year of publication, chapter title, name(s) of volume editor(s), volume title, volume number (if any), name of publisher, city of publication, and page numbers. Books should include author(s) or editor(s), year of publication, book title, publisher, and city of publication. Include only references that have been cited in the text. Provide the names of all authors; do not use "et al." in the reference list.

Examples of typical types of references follow. In addition to the particular reference styles, the examples illustrate the order in which references should be listed and give examples of "a" and "b" designations.

Bernstein, T.M., 1985. *The Careful Writer: A Modern Guide to English Usage*. Atheneum, New York.  
Buchsbbaum, M.S., 1990. Frontal lobes, basal ganglia, temporal lobes--three sites for schizophrenia? *Schizophrenia Bulletin* 16, 377-378.  
Buchsbbaum, M.S., Holcomb, H.H., DeLisi, L.E., Hazlett, E., 1986. Brain imaging in affective disorders. In: Rush, A.J., Altshuler, K.Z. (Eds.), *Depression: Basic Mechanisms, Diagnosis and Treatment*. The Guilford Press, New York, pp. 126-142.

Issa, F., Gerhardt, G.A., Bartko, J.J., Suddath, R.L., Lynch, M., Gamache, P.H., Freedman, R., Wyatt, R.J., Kirch, D.G., 1994a. A multidimensional approach to analysis of cerebrospinal fluid biogenic amines in schizophrenia: I. Comparisons with healthy control subjects and neuroleptic-treated/unmedicated pairs analyses. *Psychiatry Research* 52, 237-249.

Issa, F., Kirch, D.G., Gerhardt, G.A., Bartko, J.J., Suddath, R.L., Freedman, R., Wyatt, R.J., 1994b. A multidimensional approach to analysis of cerebrospinal fluid biogenic amines in schizophrenia: II. Correlations with psychopathology. *Psychiatry Research* 52, 251-258.

Strunk, W., White, E.B., 1979. *The Elements of Style*, 3rd ed. MacMillan, New York.

The correctness of the reference list is the entire responsibility of the author! Please check it carefully and remember to recheck when your article has been revised. Unpublished results should not be included in the reference list but, rather, should be quoted in the text (Smith and co-workers, unpublished results).

#### *Reference links*

Increased discoverability of research and high quality peer review are ensured by online links to the sources cited. In order to allow us to create links to abstracting and indexing services, such as Scopus, CrossRef and PubMed, please ensure that data provided in the references are correct. Please note that incorrect surnames, journal/book titles, publication year and pagination may prevent link creation. When copying references, please be careful as they may already contain errors. Use of the DOI is encouraged.

#### *Web references*

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

#### *Journal abbreviations source*

Journal names should be abbreviated according to the List of Title Word Abbreviations: <http://www.issn.org/services/online-services/access-to-the-ltwa/>.

#### **Video data**

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the files in one of our recommended file formats with a preferred maximum size of 50 MB. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our video instruction pages at <http://www.elsevier.com/artworkinstructions>. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

#### **AudioSlides**

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. More information and examples are available at <http://www.elsevier.com/audioslides>. Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

#### **Supplementary data**

Elsevier accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high-resolution images, background datasets, sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. In order to ensure that your submitted material is

directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. For more detailed instructions please visit our artwork instruction pages at <http://www.elsevier.com/artworkinstructions>.

### **3D neuroimaging**

You can enrich your online articles by providing 3D neuroimaging data in NIFTI format. This will be visualized for readers using the interactive viewer embedded within your article, and will enable them to: browse through available neuroimaging datasets; zoom, rotate and pan the 3D brain reconstruction; cut through the volume; change opacity and color mapping; switch between 3D and 2D projected views; and download the data. The viewer supports both single (.nii) and dual (.hdr and .img) NIFTI file formats. Recommended size of a single uncompressed dataset is 100 MB or less. Multiple datasets can be submitted. Each dataset will have to be zipped and uploaded to the online submission system via the '3D neuroimaging data' submission category. Please provide a short informative description for each dataset by filling in the 'Description' field when uploading a dataset. Note: all datasets will be available for downloading from the online article on ScienceDirect. If you have concerns about your data being downloadable, please provide a video instead. For more information see: <http://www.elsevier.com/3DNeuroimaging>.

### **Brain images and statistical images**

Brain images should be of sufficient size that small marked regions are clearly visible. Large areas of surrounding black background should be minimized so that the brain images make up the majority of the illustration. Enlargements of portions of the brain should be considered as an additional part of the illustration.

In color illustrations please provide a color bar with scale points marked so that the colors will be interpretable.

The legend to the figure should mention the right-left orientation of the figure.

### **Submission checklist**

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

#### **Ensure that the following items are present:**

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address
- Phone numbers

All necessary files have been uploaded, and contain:

- Keywords
- All figure captions
- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
- References are in the correct format for this journal
- All references mentioned in the Reference list are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)

Printed version of figures (if applicable) in color or black-and-white

- Indicate clearly whether or not color or black-and-white in print is required.
- For reproduction in black-and-white, please supply black-and-white versions of the figures for printing purposes.

For any further information please visit our customer support site at <http://support.elsevier.com>.

### **AFTER ACCEPTANCE**

#### **Use of the Digital Object Identifier**

The Digital Object Identifier (DOI) may be used to cite and link to electronic documents. The DOI consists of a unique alpha-numeric character string which is assigned to a document by the publisher upon the initial electronic publication. The assigned DOI never changes. Therefore, it is an ideal

medium for citing a document, particularly 'Articles in press' because they have not yet received their full bibliographic information. Example of a correctly given DOI (in URL format; here an article in the journal *Physics Letters B*):

<http://dx.doi.org/10.1016/j.physletb.2010.09.059>

When you use a DOI to create links to documents on the web, the DOIs are guaranteed never to change.

#### **Online proof correction**

Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

#### **Offprints**

The corresponding author, at no cost, will be provided with a personalized link providing 50 days free access to the final published version of the article on [ScienceDirect](#). This link can also be used for sharing via email and social networks. For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's WebShop (<http://webshop.elsevier.com/myarticleservices/offprints>). Authors requiring printed copies of multiple articles may use Elsevier WebShop's 'Create Your Own Book' service to collate multiple articles within a single cover (<http://webshop.elsevier.com/myarticleservices/booklets>).

#### **AUTHOR INQUIRIES**

You can track your submitted article at [http://help.elsevier.com/app/answers/detail/a\\_id/89/p/8045/](http://help.elsevier.com/app/answers/detail/a_id/89/p/8045/).

You can track your accepted article at <http://www.elsevier.com/trackarticle>. You are also welcome to contact Customer Support via <http://support.elsevier.com>.

© Copyright 2014 Elsevier | <http://www.elsevier.com>